

# Evaluation of the potential value of the F1H and F2H Diatomaceous earth formulations as grain protectants against *Rhyzopertha dominica* (Fabricius) (Coleoptera: Bostrichidae)

---

Liška, Anita; Korunić, Z.; Lucić, Pavo; Baličević, Renata; Halamić, J.; Galović, I.

Source / Izvornik: **Proceedings of the 12th International Working Conference on Stored Product Protection (IWCSP), 2018, 540 - 546**

Conference paper / Rad u zborniku

Publication status / Verzija rada: **Published version / Objavljena verzija rada (izdavačev PDF)**

<https://doi.org/10.5073/jka.2018.463.118>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:151:417966>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-12-03**



Sveučilište Josipa Jurja  
Strossmayera u Osijeku

**Fakultet  
agrobiotehničkih  
znanosti Osijek**

Repository / Repozitorij:

[Repository of the Faculty of Agrobiotechnical  
Sciences Osijek - Repository of the Faculty of  
Agrobiotechnical Sciences Osijek](#)



- CHEN QIZONG, 1994. Investigation and Study on the Pests of Stored Products in China: A Discussion on the Historical Insect Pests of the National Food System[J]. *Cereals Science and Technology*, (5): 6-9.
- LI XIAODA, LI GUOCHANG, HAO LINGJUN, 1988. Investigation and Research on Stored Mites in Henan Province[J]. *Journal of Henan University of Technology(Natural Science Edition)*, (4): 67-72.
- TIAN JIANGUO, REN ZHENGONG, ZHANG TAO, 1988. Preliminary Report on the Composition and Occurrence Situation of Grain Anchovy in Shaanxi Province[J]. *Grain Processing*, (4): 3-6.
- CHEN QIZONG, 1990. Preliminary Report on the Investigation of the Fauna (Insect and Mites) in the Tibet Autonomous Region[J]. *Journal of Henan University of Technology(Natural Science Edition)*, (3): 29-41.
- WU GUOXIONG, ZHENG WEI, LAN BO, et al, 1990. Survey of stored mussels in Jiangxi Province [J]. *Grain Storage*, (4): 15-22.
- PALYVOS N E, EMMANOUEL N G, 2009. Temperature-dependent development of the predatory mite *Cheyletus malaccensis* (Acari: Cheyletidae)[J]. *Experimental & Applied Acarology*, 47(2): 147-158.
- SHEN ZHAOPENG, 1975. Life history of the first Chinese meat carp and the meat carp of *Cheyletus malaccensis* in China[J]. *ACTA ENTOMOLOGICA SINICA*, 18(3): 316-324.
- HE PEIHUAN, CAO YANG, WU YI, et al. A device for observation and storage of aphids and insects: China, 201520476487.0[P]. 2015-09-20.
- KUCEROVA Z, HROMADKOVA J, 2009. Egg Morphology of the Predatory mite, *Cheyletus malaccensis* (Acarina: Cheyletidae)[J]. *Entomologia Generalis*, 32(1):35-40.
- NAKATANI YOSHIYUKI, 1975. *Cheyletus malaccensis* Oudemans, Patterns of each stage of 1903 [J]. *Hygienic animals*, 26:151-165.
- ZHANG YANMIAO, HOU AIPING. Study on the Relationship between Meat Carp and Malaria in Malacca[J]. *Fujian Journal of Agricultural Sciences*, (1): 44-47.
- SALEH S M, ELHELALY M S, ELGAYAR F H, 1986.. Life history of the predatory mite *Cheyletus malaccensis* (Oudemans)[J]. *Acarologia*,
- PALYVOS N E, EMMANOUEL N G, 2011., Reproduction, survival, and life table parameters of the predatory mite *Cheyletus malaccensis* (Acari: Cheyletidae) at various constant temperatures[J]. *Experimental & Applied Acarology*, 54(2): 139-150.
- PULPAN J, VERNER P H, 1965. Control of Tyroglyphoid Mites in Stored Grain by the Predatory Mite *Cheyletus eruditus* (Schrank)[J]. *Canadian Journal of Zoology-revue Canadienne De Zoologie*, 43(3): 417-432.

## Evaluation of the potential value of the F<sub>1</sub>H and F<sub>2</sub>H Diatomaceous earth formulations as grain protectants against *Rhyzopertha dominica* (Fabricius) (Coleoptera: Bostrichidae)

Anita Liška<sup>1\*</sup>, Zlatko Korunić<sup>2</sup>, Vlatka Rozman<sup>1</sup>, Pavo Lucić<sup>1</sup>, Renata Baličević<sup>1</sup>, Josip Halamić<sup>3</sup>, Ines Galović<sup>3</sup>

<sup>1</sup>University of Josip Juraj Strossmayer in Osijek, Faculty of Agriculture in Osijek, Vladimira Preloga 1, 31000 Osijek, Croatia

<sup>2</sup>Diatom Research and Consulting Inc., 14 Tidefall Dr. Toronto, ON, M1W 1J2, Canada

<sup>3</sup>Croatian Geological Survey, Sachsova 2, 10000 Zagreb, Croatia

Corresponding author Email: [aliska@pfos.hr](mailto:aliska@pfos.hr)

DOI 10.5073/jka.2018.463.118

### Abstract

An insecticidal efficacy of two **newly developed** grain protectant formulations were assessed against lesser grain borer *Rhyzopertha dominica* (Fabricius) (Coleoptera: Bostrichidae) on wheat and corn after 6 months period of. Tested formulations, marked as F<sub>1</sub>H and F<sub>2</sub>H, based on inert dust, laurel leaves, lavender essential oil, corn oil, silica gel (both F<sub>1</sub>H and F<sub>2</sub>H) and pyrethrin (only F<sub>2</sub>H) were tested at six doses (from 100 ppm to 600 ppm) depending on formulation and type of grain. The appropriate weights of each formulation, were added separately to plastic containers containing 10 kg of wheat or corn. An initial population of 200 adults of *R. dominica* were added into each container and left under natural environmental conditions for up to 6 months. A commercial diatomaceous earth (DE) insecticide, Celatom<sup>®</sup> Mn 51, was used for the comparison of the results, in addition to the untreated control. After six months, both formulations showed higher insecticidal effect than DE Mn 51 in corn and in wheat. Furthermore, the initial population of *R. dominica*, introduced in wheat was suppressed almost completely, with only 0.7%-5.3% live adults found, depending on formulations and dose. The order of efficacy was F<sub>1</sub>H>F<sub>2</sub>H>DE Mn 51. Similar suppression of the initial population was recorded in corn, where F<sub>2</sub>H was slightly more effective than F<sub>1</sub>H with 2.0%-10.6% and 4.1%-9.5% live adults found, respectively. At the same time, in the treatments with DE Mn 51 there were 4.7%-74.7% and 33.4%-56.1% live adults in wheat and corn, respectively.

**Keywords:** inert dust, botanicals, grain protectant, stored product insects, insecticidal effect

## Introduction

Minimising food commodity losses, both qualitative and quantitative, during longer period of storing represents a main challenge for all economies. Stored-product insects play a significant role in postharvest losses, causing losses in grain weight, affects on baking quality and seed viability (Sánchez-Maríñez et al., 1997; Stejskal et al., 2015), which lower cereal market value. The use of synthetic insecticides is globally the most common way of controlling stored product insect pests of the negative effects of pesticides on stored products includes: toxic residues (Fang et al., 2002), resistant strains within the insect populations (Chaudhry, 2000; Boyer et al., 2012), adverse effect on human health and environment (Fields and White, 2002). Thus there is an urgent need for alternative strategies which would be sufficiently effective against insects but less toxic for the environment.

The use of inert dusts, especially diatomaceous earth (DE), suits most of those requirements. Its main advantages are low mammalian toxicity and stability (Maceljski and Korunic, 1972; Subramanyam and Roesli, 2000) and an efficient insecticidal activity without leaving hazardous residues (Korunic, 1998; Shah and Khan, 2014; Liška et al., 2015; Korunić et al., 2017). Despite this there are several limitations which hinder wider commercial use of DE for direct mixing with grains and are described by Korunić (2016). Diatomaceous earths, inert dusts in general, have physical mode of action and therefore act more slowly than conventional contact insecticides. Depending greatly on ambient conditions, it could take a several days to control most target insect species (Korunić et al, 2016), providing enough time for oviposition. Further, there are different sensitivities of insect species to DEs, varying effects of DE on insects depending upon the commodity being treated and a negative effect on bulk density (Korunic, 2016; Korunić et al, 2017). Possible solutions for minimising or avoiding those implications include incorporating DE with other methods, such as extreme temperature (Dowdy, 1999), mixture with synthetic insecticides (Athanasios, 2006; Korunic and Rozman, 2010), mixture with entomopathogenic fungi (Batta and Kavallieratos, 2018) or with botanicals (Korunic et al, 2014; Adarkwah et al, 2017).

Most experiments with inert dusts and their mixtures are carried out in controlled conditions, and less in the real conditions where various environmental and storage conditions could impact efficacy and subsequently, storage duration of the commodity.

The objective of this study was to test insect activity of two new developed formulations based on Croatian inert dust, bay leaves, lavender essential oil, bait, corn oil, silica gel and pyrethrin against *R. dominica* in wheat and corn grain after six months period of storage.

## Materials and Methods

### Test insects

A local strain of *R. dominica*, was used in the experiments. Insects were reared on clean soft whole wheat kernels under controlled conditions (28±2°C, 65±5% RH, in dark). Two hundred, unsexed adults (7-21 days old) were used for each treatment.

### Commodity

Locally available commercial corn and soft wheat were used in the treatments. Commodities were sifted prior to tests in order to segregate broken kernels and other impurities. Grain moisture content and grain temperature were measured by the GAC 2100-Agri Grain analysis computer (Dickey-john). The initial measurements were 11.1 % m. c. and 23.2°C for wheat, and 10.7 % m. c. and 23.3°C for corn. Ten kg of clean wheat and corn grain was used for each treatment.

### Formulations

Two powder formulations labelled as F<sub>1</sub>H and F<sub>2</sub>H, were based on an inert dust of Croatian origin, dried and milled bay leaves, essential oil of lavender (Lavender x Intermedia), bait, corn oil, silica gel

and pyrethrin (only F<sub>2</sub>H). A commercial DE insecticide, Celatom<sup>®</sup> Mn 51, was used for the comparison of the results. It belongs to a group of DE's with medium to high efficacy against stored-product insects. Formulations and DE were tested at different doses depending on the treatment. In wheat, F<sub>1</sub>H and DE Celatom<sup>®</sup> Mn 51 were tested at 300, 400 and 500 ppm, and F<sub>2</sub>H at 100, 150 and 200 ppm, while in corn F<sub>1</sub>H and DE Celatom<sup>®</sup> Mn 51 were tested at 400, 500 and 600 ppm, and F<sub>2</sub>H at 200, 250 and 300 ppm.

### Bioassay

The dose rates of the tested formulations and DE were chosen based on results of preliminary laboratory test (unpublished data). Plastic containers of 15 L volume were filled with 10 kg of clean wheat or corn respectively. The appropriate weights of the formulations or Celetom DE were added into each container and mixed thoroughly with a power drill. Containers with untreated grain served as controls. After the dust had settled, 200 unsexed, 7-21 days old adults of *R. dominica* were added into each container which were then closed with perforated plastic lids. The bioassay was conducted for six months storing the containers in a wooden structure which simulated an average floor warehouse. The air temperature during the entire period of storing varied between 18.5°C and 23.0°C and relative humidity between 55.0% and 82.0%. After the end of the bioassay trial, the entire content of the plastic containers was sieved and all insects, dead and live, were counted.

### Results


After six months of storing of treated corn and wheat, tested formulations F<sub>1</sub>H and F<sub>2</sub>H showed different efficacy against *R. dominica* depending on the cereal type and the applied dose.


In wheat (Figure 1), both formulations successfully preserved grain quality against *R. dominica* during the period of six months. Even at the lowest dose, both formulations (300 ppm of F<sub>1</sub>H and 100 ppm of F<sub>2</sub>H) almost completely suppressed the initial population of *R. dominica*, therefore after six months, there was no population increase with only 0.7-3.8% (depending on dose of F<sub>1</sub>H) and 3.8-5.3% (depending on dose of F<sub>2</sub>H) of the original live adults found. For comparison, Celatom<sup>®</sup> Mn 51 at the lowest dose was not effective, resulting in a population increase (1688.5% higher than the initial population) within 74.7% live insects found. Population increases were also observed in the control treatment in wheat where number of insects from the initial 200 individuals increased up to 11817 within 93.13% live adults.

In corn (Figure 2), both formulations also showed better efficacy than Celatom<sup>®</sup> Mn 51. According to the number of adults found at the end of the testing period, formulation F<sub>2</sub>H was more effective in regard to F<sub>1</sub>H. Further, even at the lowest dose (200 ppm) no population increase was observed, while F<sub>1</sub>H resulted with 62% and 14.5% of the population increase (at the dose of 400 ppm and 600 ppm, respectively). However, in the treatment with DE Mn 51, the highest dose (600 ppm) was not efficient enough to control *R. dominica*, and the population increase was in the range from 64.5 to 107.5% (higher than initial population) (depending on dose) within average of 45.4% live adults found. Concerning control treatments, unlike in wheat no population increase was observed in corn. Due to high temperature and high moisture content of the grain during six months fungi developed intensively and the whole stock become glued mass, so insects could not survive in those conditions.

**Fig. 1** Efficacy of the formulations F<sub>1</sub>H and F<sub>2</sub>H against *R. dominica* after six months of wheat storing

**Fig. 2** Efficacy of the formulations F<sub>1</sub>H and F<sub>2</sub>H against *R. dominica* after six months of corn storing





## Discussion

Increased efficacy of the tested formulations against *R. dominica*, compared to DE alone was expressed because of the mixture of different active ingredients within their composition and due to their different modes of action. Besides inert dusts, the main composition of the formulations are the botanicals powdered bay leaves and lavandin essential oil. These compounds possess fast toxic activity against many coleopteran pests of stored products (Kostyukovsky et al 2002; Rajendran and Sriranjini, 2008; Nerio et al 2010; Koutsaviti et al, 2018). Comprised of volatile monoterpenoids and sesquiterpenoids, the essential oils interfere with basic metabolic, biochemical, physiological and behavioural functions in insects possessing contact, inhalation and ingestion toxicity, antifeedant activity, developmental delay of adult emergence and fertility, different effects on oviposition and repellent activity (Obeng-Ofori 2007; Caballero-Gallardo et al, 2012; Nenaah 2014; Germinara et al, 2017). DEs and general inert dusts, with physical mode of action, are slow acting protectants (Korunic, 1998). Apparently, as a mixture of plant powders, essential oils and inert dusts, our formulations accelerate the knock down effect of the adults within the initial population of *R. dominica* which resulted in prevention of mating and reduced oviposition. Food grade bait composed within our formulation probably also accelerated insect mortality. Presumably, it attracted insects which kept them in the contact with inert dust for longer period. Consequently, insects picked up more inert dust particles on their body which led to faster desiccation (Korunić et al, 2016).

Adarkwah (2017, 2017a) reported faster activity of the mixture of DE and plant powders against three different stored product insects. While, mentioned authors conducted their trials in laboratory conditions and only during 7 days of exposure, our tested formulations secured effective control of the tested pest during the whole testing period of six months in conditions that might more realistically represent true storage conditions.

Overall two developed formulations, as a combination of inert dusts and botanicals could be promising insecticides with residual effect for protecting stored wheat and corn against insect infestation.

## Acknowledgement

Financial support for this research was provided by the Croatian Science Foundation through scientific research project IP-11-2013-5570: "Development of new natural insecticide formulations

based on inert dusts and botanicals to replace synthetic, conventional insecticides”, [www.diacromixpest.eu](http://www.diacromixpest.eu).

## References

- ADARKWAH, C., OBENG-OFORI, D., HÖRMANN, V., ULRICHS, C. UND M. SCHÖLLER, 2017: Bioefficacy of enhanced diatomaceous earth and botanical powders on the mortality and progeny production of *Acanthoscelides obtectus* (Coleoptera: Chrysomelidae), *Sitophilus granaries* (Coleoptera: Dryophthoridae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae) in stored grain cereals. *International Journal of Tropical Insect Science*, **37** (4): 243-258.
- ADARKWAH, C., OBENG-OFORI, D., ULRICHS, C. UND M. SCHÖLLER, 2017a: Insecticidal efficacy of botanical food by-products against selected stored-grain beetles by the combined action with modified diatomaceous earth. *Journal of Plant Diseases and Protection*, **124** (3): 255-267.
- ATHANASSIOU, C.G., 2006: Toxicity of beta cyfluthrin applied alone or in combination with diatomaceous earth against adults of *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae) and *Tribolium confusum* (DuVal.) (Coleoptera: Tenebrionidae) on stored wheat. *Crop protection*, **25**: 788-794.
- BATTA, Y.A. UND N.G. KAVALLERATOS, 2018: The use of entomopathogenic fungi for the control of stored-grain insects. *International Journal of Pest Management*, **64** (1): 77-87.
- BOYER, S., ZHANGAND, H. UND G. LEMPÉRIÈRE, 2012: A review of control methods and resistance mechanisms in stored-product insects. *Bull Entomol Res*, **102** (2): 213-229.
- CABALLERO-GALLARDO, K., OLIVERO-VERBEL, J. UND E.E. STASHENKO, 2012: Repellency and toxicity of essential oils from *Cymbopogon martinii*, *Cymbopogon flexuosus* and *Lippia organoides* cultivated in Colombia against *Tribolium castaneum*. *Journal of Stored Products Research*, **50**: 62-65.
- CHAUDHRY, M.Q., 2000: Phosphine resistance: a growing threat to an ideal fumigant. *Pesticide Outlook*, **6**: 88-91.
- DOWDY, A.K., 1999: Mortality of red flour beetle, *Tribolium castaneum* (Coleoptera: Tenebrionidae) exposed to high temperature and diatomaceous earth combinations. *Journal of Stored Products Research*, **35**: 175-182.
- FANG, L., SUBRAMANYAM, B. UND S. DOLDER, 2002: Persistence and efficacy of spinosad residues in farm stored wheat. *Journal of Economic Entomology*, **95**: 1102-1109.
- FIELDS, P.G. UND N.D. WHITE, 2002: Alternatives to methyl bromide treatments for stored-product and quarantine insects. *Annu Rev Entomol*, **47** (1): 331-359.
- GERMINARA, G.C., DE STEFANO, M.G., DE ACUTIS, L., PATI, S., DELFINE, S.; DE CHRISTFARD, A. UND G. ROTUNO, 2017: Bioactivities of *Lavandula angustifolia* essential oil against the stored pest *Sitophilus granarius*. *Bulletin of Insectology*, **70** (1): 129-138.
- KORUNIC, Z., 1998: Review Diatomaceous Earth – A group of natural insecticides. *Journal of Stored Products Research*, **34**: 87-97.
- KORUNIC, Z., 2016: Overview of undesirable effects of using diatomaceous earths for direct mixing with grains. *Pectic. Phytomed. (Belgrade)*, **31** (1-2): 9-18.
- KORUNIC, Z. UND V. ROZMAN, 2010: A synergistic mixture of diatomaceous earth and deltamethrin to control stored grain insects. In: M.O. Carvalho, P.G. Fields, C.S. Adler, F.H. Arthur, C.G. Athanassiou, J.F. Campbell, et al. (Eds.), *Proceedings of the 10<sup>th</sup> International Conference on stored product protection*, Estoril, **425**: 894-898.
- KORUNIC, Z., ALMASI, R., ANDRIC, G., KLJAJIC, P., FIELDS, P.G., WAKIL, W. UND M. ZIAEE, 2014: Variation in the susceptibility of stored-product insects from five locations to two diatomaceous earth and botanical based formulations. In: F.H Arthur, R. Kengkanpanich; W. Chayaprasert, D. Suthisut (Eds.), *Proceedings of the 11<sup>th</sup> International Conference on stored product protection*, Chiang Mai, Thailand, **426**: 808-818.
- KORUNIC, Z., LIŠKA, A., ROZMAN, V. UND P. LUCIĆ, 2016: A review of natural insecticides based on diatomaceous earth. *Poljoprivreda* **22** (1): 10-18. DOI: 10.18047/poljo.22.1.2.
- KORUNIC, Z., LIŠKA, A., ROZMAN, V. UND P. LUCIĆ, 2017: Laboratory tests on insecticidal effectiveness of disodium octaborate tetrahydrate, diatomaceous earth and amorphous silica gel against *Sitophilus oryzae* (L.) and their effect on wheat bulk density. *Poljoprivreda* **23** (1): 3-10. DOI: 10.18047/poljo.23.1.1.
- KOSTYUKOVSKY, M., RAFAELI, A., GILEADI, C., DEMCHENKO, N. UND E. SHAAYA, 2002: Activation of octopaminergic receptors by essential oil constituents isolated from aromatic plants: possible mode of action against insect pests. *Pest Mgt. Sci.* **58**: 1101-1106.
- KOUTSAVITI, A., ANTONOLOULOU, V., VLASSI, A., ANTONATOS, S., MICHAELAKIS, A., PAPACHRISTOS, P.D. UND O. TZAKOU, 2018: Chemical composition and fumigant activity of essential oils from six plant families against *Sitophilus oryzae* (Col: Curculionidae), *Journal of pest Science*, **91** (2): 873-886.
- LIŠKA, A., ROZMAN, V., KORUNIC, Z., HALAMIĆ, J., GALOVIĆ, I., LUCIĆ, P. UND R. BALIĆEVIĆ, 2015: The potential of Croatian diatomaceous earths as grain protectant against three stored-product insects. *Integrated Protection of Stored Products IOBC-WPRS Bulletin*, **111**, 2015: 107-113.
- MACELJSKI, M. UND Z. KORUNIC, 1972: The effectiveness against stored-product insects of inert dusts, insect pathogens, temperature and humidity. Project No. E30-MQ-1. Grabt USDA/YU No. FG-YU-130. Final report of Institute for Plant protection. Zagreb, Croatia.
- NENAAH, G. E., 2014: Chemical composition, toxicity and growth inhibitory activities of essential oil of three *Achillea* species and their nano-emulsions against *Tribolium castaneum* (Herbst). *Industrial Crops and Products*, **53**: 252-260.
- NERIO, L. S., OLIVERO-VERBEL, J. UND E. STASHENKO, 2010: Repellent activity of essential oils: a review. *Bioresour Technol* **101** (1): 372-378.
- OBENG-OFORI, D., 2007: The use of botanicals by resource poor farmers in Africa and Asia for the protection of stored agricultural products. *Stewart Postharvest Review*, **6**: 1-8.

- RAJENDRAN, S. UND V. SRIRANJINI, 2008: Plant products as fumigants for stored product insect control. *J. Stored Prod. Res.* **44**: 126-135
- SÁNCHEZ-MARÍÑEZ, R.I., CORTEZ-ROCHA, M.O., ORTEGA-DORAME, F., MORALES-VALDES, M. UND M.I. SILVEIRA, 1997: End-use quality of flour from *Rhyzopertha dominica* infested wheat. *Cereal chemistry* **74**: 481-483.
- SHAH, M.A. UND A.A. KHAN, 2014: Use of diatomaceous earth for the management of stored-product pests. *International Journal of Pest Management*, **60** (2): 100-113.
- STEJSKAL, V., HUBERT, J., AULICKY, R. UND Z. KUCEROVA, 2015: Overview of present and past and pest-associated risks in stored food and feed products: European prospective. *Journal of Stored Products Research* **64**: 122-132.
- SUBRAMANYAM, B.H. UND R. ROESLI, 2000: Inert dusts. In (B.H. Subramanyam and D.W. Hagstrum (Eds.), *Alternatives to pesticides in stored-product IPM*. Dordrecht: Kluwer Academic Publishers: 321-380.

## Olfactory host location and host preference of *Holepyris sylvanidis* (Hymenoptera: Bethyridae) and *Cephalonomia waterstoni* (Bethyridae), two natural enemies of *Tribolium* and *Cryptolestes* species

Marco Amante<sup>1\*</sup>, Agatino Russo<sup>1</sup>, Matthias Schöller<sup>2</sup>, Johannes L.M. Steidle<sup>3</sup>

<sup>1</sup> University of Catania, Dipartimento di Agricoltura, Alimentazione e Ambiente, via Santa Sofia, 100, 95123 Catania, Italy

<sup>2</sup> Biologische Beratung GmbH, Storkower Str. 55, D-10409 Berlin, Germany

<sup>3</sup> University Hohenheim, Institut für Zoologie, Fachgebiet Tierökologie 220c, Garbenstrasse 30, 70593 Stuttgart, Germany

\* Corresponding author: marcoamante@live.it

DOI 10.5073/jka.2018.463.119

### Abstract

Parasitoids can suppress populations of their host and thus play a primary role in Integrated Pest Management. In the stored product environment, stimuli deriving from plant products, damaged plant products and hosts might be important for host location by the parasitoids. We studied foraging cues in *Holepyris sylvanidis* (Hymenoptera: Bethyridae), a larval parasitoid of *Tribolium* species and *Cephalonomia waterstoni* (Bethyridae), a natural enemy of the rusty grain beetle *Cryptolestes ferrugineus* (Coleoptera: Cucujidae). Our studies in a four-chamber olfactometer revealed that the host complexes of both *Tribolium* species and different living host stages attract naive *H. sylvanidis* females, whereas no reaction was observed to uninfested substrates. The olfactory response of *C. waterstoni* was found to be strongly elicited both by chemicals emitted by the dust, adult *C. ferrugineus* and *C. ferrugineus* third and fourth instar larvae. Our findings may contribute to the development of biological control strategies of *T. castaneum*, *T. confusum* and *C. ferrugineus* with parasitoids.

**Keywords:** natural enemies, Bethyridae, stored product pests, biological control

### Introduction

The bethylid wasp *Cephalonomia waterstoni* Gahan is an external, arrhenotokous idiobiont larval ectoparasitoid. Hosts are *Cryptolestes ferrugineus* (Stephens), *C. pusillus* (Schönherr) and *C. turcicus* (Grouvelle) (Coleoptera: Cucujidae) (Finlayson, 1950a; 1950b). *C. waterstoni* is able to find hosts by recognizing residual kairomonal cues on infested substrates, similar to other parasitoids (Howard & Flinn, 1990). Hagstrum (1987) and Reichmuth et al. (2007) reported the ability of *C. waterstoni* to maintain the population of rusty grain beetles below the economic threshold.

The parasitic wasp *Holepyris sylvanidis* (Brèthes) (Hymenoptera: Bethyridae) is a larval parasitoid of *Tribolium confusum* Jacqueline du Val and *T. castaneum* (Herbst) (Coleoptera: Tenebrionidae), the economically most important stored product pests worldwide (Athanassiou et al., 2005; García et al., 2005). The host-searching behaviour of *H. sylvanidis* is influenced by the presence of host faeces, in which two compounds are thought to be responsible for the attraction: (*E*)-2-nonenal and 1-pentadecene (Fürstenau et al., 2016). The ability of *H. sylvanidis* to penetrate cracks and crevices makes it a promising natural enemy against stored-product pests. Pest larvae are often hidden under thin layers of substrate, in aeration ducts, in machines and in areas that are difficult to clean, but this wasp is able to access these critical environments.