

LACTATE DEHYDROGENASE GENE GENOTYPIZATION FOR SPECIES IDENTIFICATION IN A FISH FARM ON THE RIVER NERETVA

Margeta, Polonca; Bogut, Ivan; Ivanković, Marko; Margeta, Vladimir

Source / Izvornik: **Poljoprivreda, 2015, 21, 56 - 58**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.18047/poljo.21.1.sup.12>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:151:021548>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-02-05**



Sveučilište Josipa Jurja
Strossmayera u Osijeku

**Fakultet
agrobiotehničkih
znanosti Osijek**

Repository / Repozitorij:

[Repository of the Faculty of Agrobiotechnical
Sciences Osijek - Repository of the Faculty of
Agrobiotechnical Sciences Osijek](#)



Lactate dehydrogenase gene genotypization for species identification in a fish farm on the river Neretva

Margeta, P., Bogut, I., Ivanković, M., Margeta, V.

Poljoprivreda/Agriculture

ISSN: 1848-8080 (Online)

ISSN: 1330-7142 (Print)

<http://dx.doi.org/10.18047/poljo.21.1.sup.12>



Poljoprivredni fakultet u Osijeku, Poljoprivredni institut Osijek

Faculty of Agriculture in Osijek, Agricultural Institute Osijek

LACTATE DEHYDROGENASE GENE GENOTYPIZATION FOR SPECIES IDENTIFICATION IN A FISH FARM ON THE RIVER NERETVA

Margeta, P.⁽¹⁾, Bogut, I.⁽¹⁾, Ivanković, M.⁽²⁾, Margeta, V.⁽¹⁾

Original scientific paper

SUMMARY

*There are several Salmonid species, found in the river Neretva basin, among which *S. trutta* and *S. obtusirostris*. Also, natural hybrids such as *S. obtusirostris* x *S. trutta* have been observed. In one fish farm on the river Neretva, *S. trutta* and *S. obtusirostris* were decided to breed separately. Parental fishes were separated phenotypically on the basis of the morphological signs. PCR-RFLP analysis of the exon 3 to exon 4 part of the lactate dehydrogenase (LDH) C1* gene with restriction endonuclease *RsaI* was employed to identify the presence of other species representatives or intercrosses in two groups of juvenile fishes. Using this method, we were able to identify two *S. trutta* representatives in the *S. obtusirostris* group.*

Key-words: Lactate dehydrogenase gene, PCR-RFLP, *S. trutta*, *S. obtusirostris*

INTRODUCTION

There are several Salmonid species, found in the river Neretva basin, among which *S. trutta* (brown trout), *Salmo marmoratus* (marble trout), *Salmo obtusirostris* (softmouth trout) *Salmo farioides* and *Salmo dentex*.

Salmo obtusirostris also known as the Adriatic trout, Adriatic salmon, and softmouth trout, is a species of salmonid fish endemic to the rivers of Western Balkans. It is found naturally in four drainages of the Adriatic Sea, in Croatia, Bosnia and Herzegovina and Montenegro: the Neretva-Vrljika system, the Jadro, the Morača-Zeta system and the Krka river drainages. Morphological different characteristics for different softmouth trout populations from different river-systems resulted in the description of three putative subspecies: *Salmo obtusirostris oxyrhynchus* from the River Neretva, Bosnia and Herzegovina, *Salmo obtusirostris salonitana* from the river Jadro, Croatia, and *Salmo obtusirostris krkensis* from the River Krka, Croatia.

In the river Neretva basin, natural hybrids such as *S. obtusirostris* x *S. trutta* have been observed and reported (Vuković, 1982). The hybridization between autochthonous species was confirmed also in an experiment performed in the fish farm located in the river Buna, a tributary of the river Neretva (Kosorić and Vuković, 1969). Introduction of non-native brown trout

has also been practised in the river Neretva and stocking activities have never been well documented.

Lactate dehydrogenase (LDH) C1* gene containing parts of exons 3 and 4 with intermediate intron has been found, firstly in brown trout (Mc Meel et al, Ferguson, 2001), and after that also in *S. obtusirostris* (Snoj et al., 2002) proven as an informative genetic marker.

A mutation in the intron 3 of the (LDH) C1* gene (G/C substitution) was connected with the disruption of the *RsaI* restriction enzyme recognition site, being specific for *S. obtusirostris* (Razpet, 2004; Razpet, 2007).

The same mutation was found in *Salmo obtusirostris salonitana* and in *Salmo obtusirostris oxyrhynchus* (Ođak, 2004).

In one fish farm on the river Neretva, *S. trutta* and *S. obtusirostris* were decided to breed separately.

Parental fishes were separated phenotypically on the basis of the morphological signs. The aim of this work was, based on the LDH genotype, to find intercrosses or representatives of the other species in two groups of juvenile fishes. In the first group, there should

(1) Ph.D. Polonca Margeta (pfrajman@gmail.com), Ph.D. Ivan Bogut, Assist. Prof. Margeta Vladimir – Josip Juraj Strossmayer University of Osijek, Faculty of Agriculture in Osijek, Kralja Petra Svačića 1d, 31000 Osijek, Croatia, (2) Ph.D. Marko Ivanković - Federal Agromediterranean Institute Mostar, Biskupa Čule 10, 88000 Mostar, Bosnia and Herzegovina




Figure 2. PCR-RFLP results on agarose gel. Samples 1-11 are from the *S. trutta* group and they are all of NP genotype with tree bands after restriction. Samples 13-24 are from the *S. obtusirostris* group. While samples 13-22 possess MP genotype, specific for *S. obtusirostris*, samples 23 and 24 have NP genotype. Sample 12 is 50bp marker

In the both analyzed groups no intercrosses of MP and NP genotypes were found. The assumption is that the presence of the NP genotype in the group of juvenile softmouth trutt is either consequence of the presence of pure *S. trutta* fish in the *S. obtusirostris* parental group, or juvenile fish were accidentally mixed during handling. In the fish fry morphological signs, characteristic for adult fishes of both species, were not well defined and distinguishing between representatives of both species is difficult. Based on the results, the suggestion was to first divide adult fishes into two groups, based on morphological signs. Further, parental lines of both species should be marked and sampled for genetic analysis for detection of intercrosses or representatives of other species in parental groups. Combination of morphological signs and genetic methods was already proven to be very powerful method in distinguishing between *S. marmoratus* and hybrids with alien *Salmo* species in the Soča river basin (Delling et al., 2000).

Disadvantage of the McNeel's method is that it could be used only to elucidate *S. obtusirostris* representatives from other *Salmo* species, found in the river Neretva. While the MP genotype of the (LDH) C1* gene is a characteristic only for *S. obtusirostris*, the NP genotype is not present only in *S. trutta*, but also in *S. marmoratus* and other *Salmo* species, found in the river Neretva basin

(Razpet, 2006). Other genetic methods like microsatellite genotyping, mitochondrial DNA analysis and SNP genotyping were described in different phylogenetic studies of *Salmo* species in the river Neretva (Snoj et al., 2002; Razpet et al., 2007; Pustovrh et al., 2014). The described methods could also be used for species identification in fish farms. The problem is that these methods are much more expensive than PCR-RFLP genotyping and therefore they are financially not suitable for the fish farm owners.

CONCLUSIONS

PCR-RFLP analysis of the exon 3 to exon 4 part of the lactate dehydrogenase (LDH) C1* gene was employed to identify the presence of other species representatives or intercrosses in the two groups of salmonides (*S. trutta* and *S. obtusirostris* group), raised in a fish farm on the river Neretva. Using this method, we were able to identify two *S. trutta* representatives in the *S. obtusirostris* group. There is also a great interest of fish farmeres to raise other salmonides from the river Neretva with a help of genetic methods. Due to financial limitation, there is a need for developing the low-cost genetic methods distinguishing between other commercially interesting salmonides from the river Neretva.

REFERENCES

1. Kosorić, Đ., Vuković, T. (1969): Ispitivanja mogućnosti hibridizacije salmonidnih vrsta sliva Neretva. *Ichthyologia*, 1: 57–67.
2. Mc Meel, O.M., Hoey, E.M., Ferguson, A. (2001): Partial nucleotide sequences, and routine typing by polymerase chain reaction-restriction fragment length polymorphism, of the brown trout (*Salmo trutta*) lactate dehydrogenase, LDHC1* 90 and *100 alleles. *Molecular Ecology*, 10: 29–34.
3. Odak Tomljanović, T. (2004): Molecular-biologic characterization of endemic softmouth trout *Salmothymus obtusirostris* salonitana. Master thesis, Agronomski fakultet Sveučilišta u Zagrebu.
4. Pustovrh, G., Snoj, A., Sušnik Bajec, S. (2014): Molecular phylogeny of *Salmo* of the western Balkans, based upon multiple nuclear loci. *Genetics Selection Evolution*, 46: 7. doi: <http://dx.doi.org/10.1186/1297-9686-46-7>
5. Razpet, A., Sušnik, S., Jug, T., Snoj, A. (2007): Genetic variation among trout in the river Neretva basin, Bosnia and Herzegovina. *J. Fish. Biol.*, 70: 94–110. doi: <http://dx.doi.org/10.1111/j.1095-8649.2007.01392.x>
6. Razpet, A. (2004): Genetic diversity of salmonids of the river Neretva drainage. Master thesis, Biotehniška univerza v Ljubljani, Oddelek za zootehniko.
7. Snoj, A., Melkić, E., Sušnik, S., Muhamedagić, S., Dovč, P. (2002): DNA phylogeny supports revised classification of *Salmothymus obtusirostris*. *Biological Journal of the Linnean Society* 77: 399–411. doi: <http://dx.doi.org/10.1046/j.1095-8312.2002.00130.x>
8. Vuković, T. (1982): *Sistematika riba*. In: *Slatkovodno ribarstvo* (Habeković, D., ed.), pp. 99–168. Zagreb: Jugoslavenska medicinska naklada.

(Received on 8 June 2015; accepted on 3 August 2015)