

Alelopatsko djelovanje nekih biljnih vrsta na rast i razvoj usjeva i korova

Ravlić, Marija

Source / Izvornik: **Poljoprivreda, 2016, 22, 53 - 53**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.18047/poljo.22.1.8>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:151:544925>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-10-02**



Sveučilište Josipa Jurja
Strossmayera u Osijeku

**Fakultet
agrobiotehničkih
znanosti Osijek**

Repository / Repozitorij:

[Repository of the Faculty of Agrobiotechnical
Sciences Osijek - Repository of the Faculty of
Agrobiotechnical Sciences Osijek](#)



Alelopatsko djelovanje nekih biljnih vrsta na rast i razvoj usjeva i korova

Allelopathic effects of some plant species on growth and development of crops and weeds

Marija Ravlić

Poljoprivreda/Agriculture

ISSN: 1848-8080 (Online)

ISSN: 1330-7142 (Print)

<http://dx.doi.org/10.18047/poljo.22.1.8>



Poljoprivredni fakultet u Osijeku, Poljoprivredni institut Osijek
Faculty of Agriculture in Osijek, Agricultural Institute Osijek

ISSN 1330-7142

UDK: 581.6:633+632.5

DOI: 10.18047/poljo.22.1.8

ALELOPATSKO DJELOVANJE NEKIH BILJNIH VRSTA NA RAST I RAZVOJ USJEVA I KOROVA

Marija Ravlić, mag. ing. agr. ⁽¹⁾

Disertacija ⁽²⁾

Cilj rada bio je utvrditi alelopatski utjecaj korovnih vrsta na rast i razvoj usjeva te mogućnost primjene aromatičnih i ljekovitih biljaka u suzbijanju korova. Ispitan je utjecaj biljne mase i sjemena korovnih vrsta: poljski osjak (*Cirsium arvense*), poljski mak (*Papaver rhoeas*), bezmirisna kamilica (*Tripleurospermum inodorum*), oštrodlakavi šćir (*Amaranthus retroflexus*), crna pomoćnica (*Solanum nigrum*) i divlji sirak (*Sorghum halepense*) na pšenicu, ječam, mrkvu, soju i uljnu bundevu. Djelovanje ljupčaca (*Levisticum officinale*), bosiljka (*Ocimum basilicum*), prave kamilice (*Matricaria chamomilla*), matičnjaka (*Melissa officinalis*), crnoga sljeza (*Malva sylvestris*) i velikoga rosopasa (*Chelidonium majus*) utvrđeno je na rast i razvoj korovnih vrsta Teofrastov mračnjak (*Abutilon theophrasti*), oštrodlakavi šćir, strjeličasta grbica (*Lepidium draba*), crna pomoćnica, divlji sirak i bezmirisna kamilica. Alelopatski utjecaj ovisio je o korovnoj vrsti, usjevu, koncentraciji, odnosno dozi, biljnome dijelu te stanju biljne mase. U prosjeku su ekstrakti od svih vrsta, osim divljega sirka, smanjili klijavost za više od 20,0%, a rast klijanaca do 41,2% u Petrijevim zdjelicama. Vodeni ekstrakti primijenjeni u posude s tlom smanjili su nicanje i rast usjeva i do 65%. Inkorporacija biljnih ostataka korova u posude s tlom smanjila je nicanje usjeva i do 65,3%. Zajedničko klijanje sjemena usjeva i korova, kao i korijenovi eksudati, nisu pokazali značajan utjecaj na klijavost usjeva. Alelopatski utjecaj aromatičnoga i ljekovitoga bilja također je ovisio o biljnoj vrsti i korovu, koncentraciji i dozi te stanju biljne mase. Pri zajedničkome klijanju sjemena zabilježen je negativan utjecaj na klijanje i do 90,0%. Zajedničko klijanje sjemena u posudama s tlom slabije je djelovalo na korove. Vodeni ekstrakti od svježe i suhe mase smanjili su klijavost i rast korova i do 100%. Biljni ostatci smanjili su nicanje korovnih vrsta i do 60,0% te negativno utjecali na rast klijanaca.

Ključne riječi: alelopacija, korovi, usjevi, vodeni ekstrakti, biljni ostatci, zajedničko klijanje, eksudati

ALLELOPATHIC EFFECTS OF SOME PLANT SPECIES ON GROWTH AND DEVELOPMENT OF CROPS AND WEEDS

Doctoral thesis

The aim of the study was to determine the allelopathic effect of weeds on crop growth and development, as well as the possibility of applying aromatic and medicinal plants in weed control. The effect of biomass and seeds of weed species creeping thistle (*Cirsium arvense*), field poppy (*Papaver rhoeas*), scentless mayweed (*Tripleurospermum inodorum*), redroot pigweed (*Amaranthus retroflexus*), black nightshade (*Solanum nigrum*) and johnsongrass (*Sorghum halepense*) on wheat, barley, carrot, soybean and oil pumpkin was examined. The effect of lovage (*Levisticum officinale*), basil (*Ocimum basilicum*), chamomile (*Matricaria chamomilla*), lemon balm (*Melissa officinalis*), common mallow (*Malva sylvestris*) and greater celandine (*Chelidonium majus*) was evaluated on growth and development of weeds species velvetleaf (*Abutilon theophrasti*), redroot pigweed, hoary cress (*Lepidium draba*), black nightshade, johnsongrass and scentless mayweed. The allelopathic effect depended on the weed and crop species, concentration or rate, plant part and plant biomass. On the average, the extracts from all weed species, except johnsongrass, reduced the germination over 20.0% and growth up to 41.2% in Petri dishes. The water extracts applied in the pots reduced the crop emergence up to 65%. In the pots, the incorporation of weed residues reduced the emergence up to 65.3%. The seed cogermination and weed root exudates showed no significant effect on crop germination. The allelopathic effects of aromatic and medicinal plants depended on the plant and weed species, concentration and rate and plant biomass. The cogermination reduced the weed germination up to 90.0%. The seed cogermination showed a lower effect on weeds in the pots. The water extracts from the fresh and dry biomass reduced the germination and growth of weeds up to 100%. The plant residues reduced the weed emergence up to 60.0% and showed a negative effect on the seedling growth.

Key-words: allelopathy, weeds, crops, water extracts, plant residues, cogermination, exudates

(1) Sveučilište Josipa Jurja Strossmayera u Osijeku, Poljoprivredni fakultet u Osijeku / Josip Juraj Strossmayer University of Osijek, Faculty of Agriculture in Osijek, Kralja Petra Svačića 1d, Osijek/Croatia (mravlic@pfos.hr)

(2) Disertacija je obranjena na Sveučilištu Josipa Jurja Strossmayera, Poljoprivrednom fakultetu u Osijeku 4. prosinca 2015. godine pod mentorstvom izv. prof. dr. sc. Renate Baličević / Doctoral thesis was defended at Josip Juraj Strossmayer University of Osijek, Faculty of Agriculture in Osijek on 4th December 2015 tutored by Assoc. Prof. Renata Baličević