

STRAWBERRY (FRAGARIA X ANANASSA DUCH) LEAF ANTIOXIDATIVE RESPONSE TO BIOSTIMULATORS AND REDUCED FERTILIZATION WITH N AND K

Špoljarević, Marija; Štolfa, Ivna; Lisjak, Miroslav; Stanisavljević, Aleksandar; Vinković, Tomislav; Agić, Dejan; Parađiković, Nada; Teklić, Tihana; Engler, Meri; Klešić, Katica

Source / Izvornik: **Poljoprivreda, 2010, 16, 50 - 56**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:151:732909>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-16**



Sveučilište Josipa Jurja
Strossmayera u Osijeku

**Fakultet
agrobiotehničkih
znanosti Osijek**

Repository / Repozitorij:

[Repository of the Faculty of Agrobiotechnical
Sciences Osijek - Repository of the Faculty of
Agrobiotechnical Sciences Osijek](#)



DIGITALNI AKADEMSKI ARHIVI I REPOZITORIJI

STRAWBERRY (*FRAGARIA X ANANASSA* DUCH) LEAF ANTIOXIDATIVE RESPONSE TO BIOSTIMULATORS AND REDUCED FERTILIZATION WITH N AND K

Marija Špoljarević ⁽¹⁾, Ivna Štolfa ⁽²⁾, M. Lisjak ⁽¹⁾, A. Stanisavljević ⁽¹⁾, T. Vinković ⁽¹⁾, D. Agić ⁽¹⁾, Nada Parađiković ⁽¹⁾, Tihana Teklić ⁽¹⁾, Meri Engler ⁽¹⁾, Katica Klešić ⁽¹⁾

Original scientific paper
Izvorni znanstveni članak

SUMMARY

Strawberry cultivar Elsanta was grown in peat based substrate in a green house. Full dose and 50% reduced nitrogen and potassium fertilization were applied during fruit bearing period in spring, along with biostimulators Viva[®], Megafol[®] and their combination. The specific activities of guaiacol peroxidase (GPXs; EC 1.11.1.7), catalase (CATs; EC 1.11.1.6), ascorbate peroxidase (APXs; EC 1.11.1.11) and glutathione reductase (GRs; EC 1.6.4.2) in strawberry leaf were stimulated by biostimulators and reduced fertilization. The strongest link seen here was between the enzymes of ascorbate-glutathione cycle (APXs and GRs), which were positively related to trifoliolate leaf fresh mass (TLFM). The highest TLFM was observed in Megafol[®] treated plants.

Key-words: antioxidative activity, ascorbate peroxidase, biostimulators, catalase, fertilization, *Fragaria x ananassa* Duch, glutathione reductase, guaiacol peroxidase, leaf, strawberry

INTRODUCTION

Strawberry (*Fragaria x ananassa* Duch) is known by its attractive and delicious fruit, which has an important role in human diet because of high antioxidants content.

Strawberry is commonly grown in the field, in glasshouses, other types of green houses and plastic tunnels.

After Takeda (2000), protected cultivation systems provide opportunities to extend strawberry production to areas traditionally considered unsuitable for open-field strawberry culture. However, growth conditions in protected areas can be stressful to plants as well, particularly due higher temperature and air moisture, salt accumulation in root zone, nutrient disorders etc. All these can result with the increased production of reactive oxygen species (ROS) that are highly reactive and their reactivity dictates that they are highly energetic compounds, able to undertake catalytic functions in the absence of enzymes (Turhan et al., 2008). Tolerant

plants and genotypes can alleviate environmental stress and ROS-mediated damage through the activation of antioxidative mechanisms, among which enzymes such as different peroxidases, catalase and glutathione reductase have an important role.

Biostimulators or biostimulants are commercial liquid mixtures of different natural compounds extracted from plants, such as aminoacids, glucosides, vitamins, microelements, growth hormones and/or humic acids. There are many reports on stress-protective effects of biostimulators, followed by growth, yield and quality enhancement in different plant species (Zhang and Schmidt, 1999; Muralidharan et al., 2000; Vernieri et al., 2002; Csizinszky, 2003; Maini, 2006; Kauffman

(1) Marija Špoljarević, BSc (Marija.Spoljarevic@pfos.hr), Miroslav Lisjak, BSc, DSc Aleksandar Stanisavljević, Tomislav Vinković, BSc, Dejan Agić, BSc, Prof.DSc Nada Parađiković, Prof.DSc Tihana Teklić, Meri Engler, BSc, Katica Klešić, BSc - Faculty of Agriculture in Osijek, J.J. Strossmayer University of Osijek, Trg Sv. Trojstva 3, 31000 Osijek, Croatia; (2) Ivna Štolfa, BSc – Department of Biology, J.J. Strossmayer University of Osijek, Trg Lj. Gaja 6, 31000 Osijek, Croatia

et al., 2007; Redžepović et al., 2008; Parađiković et al., 2008, 2009). Roussos et al. (2009) observed significant positive influence of the applied mixtures of plant growth stimulators, containing seaweed extract, on strawberry yield and some fruit quality indices. However, the understanding of the physiological effects of biostimulators at the molecular level is poor. Among many different abiotic stresses that were the subject in a plethora of reports so far, nutrient supply disorders and their triggering of an oxidative stress in plants were rarely investigated, especially in the case of strawberry.

Based on the above, the aim of this research was to investigate the strawberry leaf antioxidative response to different N and K supply as well as biostimulators application in fruitbearing period. Antioxidative activity in leaf was expressed by the specific activities of guaiacol peroxidase (EC 1.11.1.7), catalase (EC 1.11.1.6), ascorbate peroxidase (EC 1.11.1.11) and glutathione reductase (EC 1.6.4.2), while the evaluated vegetative growth parameters were plant total vegetative above-ground mass, as well as fresh and dry mass of fully developed trifoliolate leaf.

MATERIAL AND METHODS

The cold stored frigo plantlets of strawberry (*Fragaria x ananassa* Duch, cultivar Elsanta) were planted in pots of 4.5 L volume, filled with commercial Stender B 400 substrate. A greenhouse was installed in the locality Dalj in the eastern Croatia. Between planting and treatment application, the plants have received 1.760 g N, 0.938 g P, 3.255 g K, 0.669 g Ca, 0.07144 g Mg and 0.189 g S per plant, as well as 6808 µg Fe, 3481 µg Mn, 1346 µg B, 1209 µg Zn, 774 µg Cu and 452 µg Mo. The experiment was set according to split plot method in 4 repetitions with 10 pots (40 plants per repetition). Reduced fertilization treatments and biostimulators were applied in the fruit bearing stage during three weeks (May 12 – June 2 2009). In fertilization variant 100% (F100) 0.540 g N, 0.973 g K, 0.453 g Ca per plant was applied, and in the reduced fertilization variant (F50) plants received 50% N and K less. Biostimulator treatments included the application of 0.25% v/v biostimulator solution of 250 mL per pot in volume. In one treatment biostimulator Viva® (B: V) was used and applied by hosing, in second Megafol® (B: M) applied by spraying, and in the third a combination of the two (B: V+M). Control plants (B: O), received 250 mL of tap water per pot. The applied biostimulators Viva® and Megafol® contain polysaccharides, glycosides, proteins, aminoacids, vitamins and essential microelements.

Leaf samples for analyses of specific enzymatic activity consisted of ten middle leaflets from the most developed leaves (triplets) per repetition, which were kept in freezer at -20°C until the analyses. Guaiacol

peroxidase activity was determined after Siegel and Galston (1967). Catalase activity was determined according to Aebi (1984), and the activity of ascorbate peroxidase was determined according to Nakano and Asada (1981). Activity of glutathione reductase was determined as described in Glutathione reductase assay kit from Oxford Biomedical Research (2001). Specific activity of analyzed enzymes is expressed in U mg⁻¹ protein, where the leaf protein concentration was determined after Bradford (1976), using bovine serum albumine (BSA) as standard.

Fresh and dry trifoliolate leaf mass (without petioles) was determined based on the ten most developed leaves taken from every repetition, before applying treatments and the week after the last application of biostimulators. Total vegetative over ground mass of plant was determined at the end of the experiment. Fresh trifoliolate leaf mass was weighed immediately, while dry mass was measured after drying at 70°C for 24 h and 2 h at 105°C.

The results were analyzed using SAS software (SAS Institute, Cary, N.C.). Analyses of variance were performed using PROC GLM with LSD test at P≤0.05. Significance of linear correlations among the tested parameters was estimated with t- test.

RESULTS AND DISCUSSION

The antioxidative enzymatic activities in strawberry leaf were significantly higher after treatments with biostimulators and reduced fertilization as compared to the same plants before treatments, except in the case of CATs activity (Table 1). As for the fertilization level influence, GPXs and CATs activities were enhanced by lower N and K supply (Table 1; P≤0.05), regarding both sampling terms (before and after the treatments) as well as if only plants after receiving treatments are considered (Table 2). In the research of Kandlbinder et al. (2004), N deprivation caused a severe drop in CAT and APX activities in photosynthesizing leaves of *Arabidopsis thaliana*. After Cakmak (2005), the need for protection against photooxidative damage in N-deficient plants can be more marked when the N deficiency stress is combined with an environmental stress. This could explain higher activities of GPXs and CATs in the leaves of plants supplied with reduced fertilization level, implying less than adequate N and K supply in that treatment. In the research of Kim et al. (2010), a member of the type III peroxidase family, RCI3, was found to be up-regulated upon K deprivation and the authors concluded that this peroxidase appears to be another component of the low-K signal transduction pathway in *Arabidopsis* roots. Shin and Schachtman (2004) reported a significant increase in ROS accumulation in plants grown under K⁺-deficient conditions and stated that the specificity of the ROS-induced responses to K deprivation are not known.

Table 1. Mean values of guaiacol peroxidase (GPX_s), catalase (CAT_s), ascorbate peroxidase (APX_s) and glutathione reductase (GR_s) specific activity in strawberry leaf under influence of term of sampling (A), fertilization level (B) and biostimulator treatment (C). (ANOVA, F test; means designated with the same letter (A,B,C) are not statistically different per protected least significant difference procedure LSD; P=0.05)

Tablica 1. Prosječne specifične aktivnosti gvajakol peroksidaze (GPX_s), katalaze (CAT_s), askorbat peroksidaze (APX_s) i glutation reduktaze (GR_s) u listu jagoda pod utjecajem termina analize (A), razine gnojidbe (B) i tretmana biostimulatorima (C). (ANOVA, F test; prosjeci označeni istim slovom (A,B,C) se ne razlikuju prema LSD testu; P=0,05)

		GPX _s	CAT _s	APX _s	GR _s
		ΔA min ⁻¹ mg ⁻¹ prot.			
Term of analyses (A) <i>Termin analize (A)</i>					
Before treatments (A1) <i>Prije tretmana (A1)</i>		1.098 ^B	5.093 ^A	1.325 ^B	1.335 ^B
After treatments (A2) <i>Poslije tretmana (A2)</i>		1.473 ^A	4.702 ^A	2.450 ^A	3.319 ^A
F test		47.36	2.25	88.81	190.36
P		<0.0001	0.1679	<0.0001	<0.0001
Fertilization variant (B) <i>Varijanta gnojidbe (B)</i>					
100% (B1)		1.199 ^B	4.489 ^B	1.796 ^A	2.203 ^A
50% (B2)		1.372 ^A	5.306 ^A	1.979 ^A	2.451 ^A
F test		10.11	9.81	2.35	2.98
P		0.0112	0.0121	0.1600	0.1186
Biostimulator treatment (C) <i>Tretman biostimulatorima (C)</i>					
Control (C1)		1.209 ^A	4.706 ^A	1.569 ^B	1.721 ^C
Viva® (C2)		1.292 ^A	4.811 ^A	1.856 ^B	2.388 ^B
Megafol® (C3)		1.291 ^A	4.87 ^A	2.407 ^A	2.953 ^A
Viva®+Megafol® (C4)		1.350 ^A	5.203 ^A	1.719 ^B	2.246 ^B
F test		1.13	0.68	9.36	12.41
P		0.3889	0.5883	0.0040	0.0015
Interactions <i>Interakcije</i>					
(AxB)	F test	10.15	6.07	7.02	4.02
P		0.0111	0.0359	0.0265	0.0759
(AxC)	F test	2.00	2.94	10.23	15.73
P		0.1848	0.0916	0.0029	0.0006
(BxC)	F test	2.91	2.22	3.51	1.38
P		0.0931	0.1547	0.0626	0.3091
(AxBxC)	F test	2.58	1.64	0.98	0.48
P		0.1187	0.2477	0.4448	0.7014

Upon changes in tissue K concentrations, it is likely that the activity of certain enzymes requiring K may be reduced (Schachtman and Shin, 2007); hence, biostimulators containing K, such as Megafol, might stimulate plant metabolism and growth through the activation of particular enzymes. Biostimulators effect on the antioxidative activity was significant in the case of APXs and GRs, considering both terms of analyses (Table 1).

Table 2. Mean values of guaiacol peroxidase (GPX_s), catalase (CAT_s), ascorbate peroxidase (APX_s) and glutathione reductase (GR_s) activity in strawberry leaf after the application of different fertilization level (A) and biostimulator treatment (B). (ANOVA, F test; means designated with the same letter (A,B,C) are not statistically different per a protected least significant difference procedure LSD; P=0.05)

Tablica 2. Prosječne vrijednosti specifične aktivnosti gvajakol peroksidaze (GPX_s), katalaze (CAT_s), askorbat peroksidaze (APX_s) i glutation reduktaze (GR_s) u listu jagoda pod utjecajem varijante gnojidbe (A) i tretmana biostimulatorima (B) nakon primjene tretmana. (ANOVA, F test; prosjeci označeni istim slovom (A,B,C) se ne razlikuju prema LSD testu; P=0,05)

		GPX _s	CAT _s	APX _s	GR _s
		ΔA min ⁻¹ mg ⁻¹ prot.			
Fertilization variant (A) <i>Varijanta gnojidbe (A)</i>					
100% (A1)		1.300 ^B	3.972 ^B	2.201 ^A	3.051 ^A
50% (A2)		1.647 ^A	5.432 ^A	2.700 ^A	3.588 ^A
F test		19.04	35.69	4.17	3.69
P		0.0018	0.0002	0.0715	0.0869
Biostimulator treatment (B) <i>Tretman biostimulatorima (B)</i>					
Control (B1)		1.325 ^A	4.161 ^B	1.706 ^B	1.973 ^C
Viva® (B2)		1.456 ^{A,B}	4.451 ^B	2.338 ^B	3.436 ^B
Megafol® (B3)		1.465 ^{A,B}	4.535 ^B	3.469 ^A	4.597 ^A
Viva®+Megafol® (B4)		1.647 ^A	5.663 ^A	2.288 ^B	3.271 ^B
F test		2.76	7.30	9.10	14.79
P		0.1039	0.0088	0.0044	0.0008
Interaction (AxB) <i>Interakcija (AxB)</i>					
F test		4.70	8.17	1.78	0.83
P		0.0308	0.0062	0.2203	0.5092

In the case of GPXs and CATs, there was a significant influence of the interaction fertilization level X biostimulator treatment. These enzymes were stimulated with lower nutrient supply (N and K; F50%) and biostimulators application (Figures 1 and

2) as well, with significant positive linear correlation established in plants after treatment ($r=0.785^*$; Table 3).




Figure 1. Guaiacol peroxidase specific activity (GPX_s) in leaf of soilless grown strawberry in green house, before and after the treatments with two-levels fertilization with N and K (F100% and F50%) and biostimulators Viva® (V) and Megafol® (M) (untreated control, Viva®, Megafol®, Viva® + Megafol®). Bars are means of four replicates with 40 plants, with standard errors ($P \leq 0.05$)

Grafikon 1. Specifična aktivnost gvajakol peroksidaze u listu jagode pri uzgoju bez tla u plasteniku, prije i nakon tretmana s dvije razine gnojidbe s N i K (F100% i F50%) te biostimulatorima Viva® (V) i Megafol® (M) (kontrola, Viva®, Megafol®, kombinacija Viva® + Megafol®). Stupci su prosječne vrijednosti četiri ponavljanja s 40 biljaka, uz standardne pogreške ($P \leq 0,05$)




Figure 2. Catalase specific activity (CAT_s) in leaf of soilless grown strawberry in green house, before and after the treatments with two-levels fertilization with N and K (F100% and F50%) and biostimulators Viva® (V) and Megafol® (M) (untreated control, Viva®, Megafol®, Viva® + Megafol®). Bars are means of four replicates with 40 plants, with standard errors ($P \leq 0.05$)

Grafikon 2. Specifična aktivnost katalaze u listu jagode pri uzgoju bez tla u plasteniku, prije i nakon tretmana s dvije razine gnojidbe s N i K (F100% i F50%) te biostimulatorima Viva® (V)

i Megafol® (M) (kontrola, Viva®, Megafol®, kombinacija Viva® + Megafol®). Stupci su prosječne vrijednosti četiri ponavljanja s 40 biljaka, uz standardne pogreške ($P \leq 0,05$)

APXs and GRs showed the highest activities in plants treated with Megafol. Taking into account only data obtained with plants after treatments, there was a significant stimulative effect of biostimulators on CATs, APXs and GRs activities. In plants before treatments, GPXs and APXs were significantly correlated to GRs (Table 3). The strongest link seen here was between the enzymes of ascorbate-glutathione cycle (APXs and GRs), which were in positive relationship regardless of term of sampling.

Table 3. Coefficients of linear correlations among the antioxidative activity parameters in strawberry leaf (guaiacol peroxidase GPX_s, catalase CAT_s, ascorbate peroxidase APX_s and glutathione reductase GR_s specific activities; * $P \leq 0,05$; ** $P \leq 0,01$)

Tablica 3. Koeficijenti linearnih korelacija između pokazatelja antioksidativne aktivnosti u listu jagode (specifične aktivnosti gvajakol peroksidaze GPX_s, katalaze CAT_s, askorbat peroksidaze APX_s i glutation reduktaze GR_s; * $P \leq 0,05$; ** $P \leq 0,01$)

Analiza prije tretmana (n=8) Analyses before treatments (n=8)		Analiza poslije tretmana (n=8) Analyses after treatments (n=8)		Oba termina analize (n=16) Both terms of analysis (n=16)	
x:y	r	x:y	r	x:y	r
GPXs:GRs	0.738*	GPXs:CATs	0.785*	GPXs:APXs	0.720**
APXs:GRs	0.789*	APXs:GRs	0.943**	GPXs:GRs	0.741**
				APXs:GRs	0.963**

While CATs activity was the highest in plants treated with the combination of two biostimulators (B4:V+M), especially in reduced fertilization treatment (Figure 2), APXs and GRs showed the highest activities in Megafol-treated plants (Figures 3 and 4), with stronger effect seen in F 50 variant. Here, the vegetative growth parameters (mean fresh trifoliolate leaf mass - TLFM; mean dry trifoliolate leaf mass - TLFM; total vegetative over ground mass per plant - TVFM) were not influenced by the applied fertilization level (Table 4). However, biostimulators effect was highly significant for TLFM ($P=0.047$; Table 4). Megafol treatment resulted in the highest values of leaf fresh and dry mass, as well as the greatest total vegetative over ground mass per plant. In the case of leaf, this growth promoting effect of Megafol was statistically significant after LSD test.




Figure 3. Ascorbate peroxidase specific activity (APX_s) in leaf of soilless grown strawberry in green house, before and after the treatments with two-levels fertilization with N and K (F100% and F50%) and biostimulators Viva® (V) and Megafol® (M) (untreated control, Viva®, Megafol®, Viva® + Megafol®). Bars are means of four replicates with 40 plants, with standard errors ($P \leq 0.05$)

Grafikon 3. Specifična aktivnost askorbat peroksidaze (APX_s) u listu jagode pri uzgoju bez tla u plasteniku, prije i nakon tretmana s dvije razine gnojidbe s N i K (F100% i F50%) te biostimulatorima Viva® (V) i Megafol® (M) (kontrola, Viva®, Megafol®, kombinacija Viva® + Megafol®). Stupci su prosječne vrijednosti četiri ponavljanja s 40 biljaka, uz standardne pogreške ($P \leq 0,05$)




Figure 4. Glutathione reductase specific activity (GR_s) in the leaf of soilless grown strawberry in green house, before and after the treatments with two-levels fertilization with N and K (F100% and F50%) and biostimulators Viva® (V) and Megafol® (M) (untreated control, Viva®, Megafol®, Viva® + Megafol®). Bars are means of four replicates with 40 plants, with standard errors ($P \leq 0.05$)

Grafikon 4. Specifična aktivnost glutatjon reduktaze (GR_s) u listu jagode pri uzgoju bez tla u plasteniku, prije i nakon tretmana s dvije razine gnojidbe s N i K (F100% i F50%) te biostimulatorima Viva® (V) i Megafol® (M) (kontrola, Viva®, Megafol®, kombinacija Viva® + Megafol®). Stupci su prosječne vrijednosti četiri ponavljanja s 40 biljaka, uz standardne pogreške ($P \leq 0,05$)

Considering that the same biostimulator showed the strongest stimulative effect on APXs and GRs activities in leaf (Table 2, Figures 3 and 4), it seems that its components (high content of amino acids and K) have certain role in ascorbate-glutathione cycle in strawberry leaf. TLFM was also influenced by the interaction fertilization level X biostimulator treatment (Table 4). The analyzed biostimulators induced statistically highly significant differences in plant height, number of branches and fresh plant mass of rosemary (*Rosmarinus officinalis* L) seedlings in the research of Jelačić et al. (2007).

Table 4. Strawberry vegetative growth parameters (mean fresh trifoliolate leaf mass - TLFM; mean dry trifoliolate leaf mass - TLDM; total vegetative over ground mass per plant - TVFM) after the application of different fertilization level (A) and biostimulator treatment (B). (ANOVA, F test; means designated with the same letter (A,B,C) are not statistically different per a protected least significant difference procedure LSD; $P = 0.05$)

Tablica 4. Pokazatelji vegetativnog rasta jagoda (prosječna svježa masa troliske - TLFM; prosječna suha masa troliske - TLDM; ukupna vegetativna masa po biljci - TVFM) nakon primjene različite gnojidbe (A) i tretmana biostimulatorima (B). (ANOVA, F test; prosjeci označeni istim slovom (A,B,C) se ne razlikuju prema LSD testu; $P = 0,05$)

	TLFM	TLDM	TVFM
	g	g	g plant ⁻¹
Fertilization variant (A) Varijanta gnojidbe (A)			
100 % (A1)	2.406 ^A	0.843 ^A	45.544 ^A
50 % (A2)	2.450 ^A	0.868 ^A	48.925 ^A
F test	0.31	0.43	4.02
P	0.5916	0.5276	0.0758
Biostimulator treatment (B) Tretman biostimulatorima (B)			
Control (B1)	2.338 ^{B,C}	0.806 ^{A,B}	46.525 ^A
Viva® (B2)	2.500 ^{A,B}	0.904 ^{A,B}	46.575 ^A
Megafol® (B3)	2.713 ^A	0.926 ^A	49.600 ^A
Viva® + Megafol® (B4)	2.163 ^C	0.784 ^B	46.238 ^A
F test	8.88	3.43	0.88
P	0.0047	0.0656	0.4858
Interaction (AxB) Interakcija (AxB)			
F test	6.52	1.48	3.20
P	0.0123	0.2858	0.0766

Considering the similar effect of Megafol on mentioned enzymes in 100% N and K fertilization level as observed here (Figures 3 and 4), it can be assumed that an explanation of positive effect of Megafol on enzyme activities and leaf fresh mass is not based on simple additional nutrient supply (N and K) by this biostimulator

to plants. However, the usage of aminoacids in biostimulators as an easy available, low molecular, organic N source for plant can not be neglected.

CONCLUSION

GPXs and CATs activities were enhanced by lower N and K supply regarding both sampling terms as well as if only plants after treatments are considered and there was a significant influence of the interaction fertilization level X biostimulator treatment. These enzymes were stimulated with lower nutrient supply and biostimulators application. Megafol treatment resulted in the highest values of leaf fresh and dry mass, as well as the greatest total vegetative over ground mass per plant. The strongest link seen here was between the enzymes of ascorbate-glutathione cycle (APXs and GRs), which were in positive relationship regardless the term of sampling. In conclusion, the understanding of feasibly complex effects of various biostimulators on plants growth, stress tolerance and productivity, imply further and more comprehensive research.

ACKNOWLEDGEMENTS

This work was an integral part of the research project no.: 079-0790494-0559 ("Physiological mechanisms of plant tolerance to abiotic stress") supported by The Ministry of science, education and sports, Croatia. We thank the administration, employees and students of the high school Dalj (<http://www.ss-dalj.skole.hr/skola>) on a chance to set up the experiment in their green house, as well as for much appreciated help during the research.

REFERENCES

- Aebi, H. (1984): Catalase *in vitro*. *Methods in Enzymology* 105: 121-126.
- Bradford, M.M. (1976): A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72: 248-254.
- Cakmak, I. (2005): Role of mineral nutrients in tolerance of crop plants to environmental stress factors. In: *Fertigation: Optimizing the utilization of water and nutrients. Fertigation Proceedings: Selected papers of the IPI-NATESC-CAU-CAAS. International Symposium on Fertigation, Beijing, China, 20-24 September 2005: 35-48.*
- Csizinszky, A.A. (2003): Response of „Florida 47“ tomato to soil and foliar-applied biostimulants and N and K rates. 116. Annual Meeting of the Florida State Horticultural Society. Program and abstract book, pp 125.
- Jelačić, S., Beatović, D., Lakić, N., Vujošević, A. (2007): The effect of natural biostimulators and slow-disintegrating fertilizers on the quality of rosemary seedlings (*Rosmarinus officinalis* L.). *Journal of Agricultural Sciences* Vol. 52, No 2, 85-94
- Kandlbinder, A., Finkemeier, I., Wormuth, D., Hanitzsch M., Dietz, K.J. (2004): The antioxidant status of photosynthesizing leaves under nutrient deficiency: redox regulation, gene expression and antioxidant activity in *Arabidopsis thaliana*. *Physiologia Plantarum* Volume 120 Issue 1, 63-73.
- Kauffman III, G.L., Kneivel, D. P., Watschke, T.L. (2007): Effects of a biostimulant on the heat tolerance associated with photosynthetic capacity, membrane thermostability, and polyphenol production of perennial ryegrass. *Crop Science* 47: 261-267.
- Kim, M.J., Ciani, S., Schachtman, D.P. (2010): A peroxidase contributes to ROS production during *Arabidopsis* root response to potassium deficiency. *Molecular Plant*, doi:10.1093/mp/ssp121; *Molecular Plant Advance Access* published online on February 5, 2010.
- Maini, P. (2006): The experience of the first biostimulant, based on aminoacids and peptides: a short retrospective review on the laboratory researches and the practical results. Ed. Centro Scientifico Italiano dei Fertilizzanti, *Fertilitas Agrorum*, 1(1): 29-43.
- Muralidharan, R., Saravanan, A., Muthuvel, P. (2000): Influence of biostimulants on yield and quality of tomato (*Lycopersicon esculentum* Mill.). *Madras Agricultural Journal*, 87 (10/12): 625-628.
- Nakano, Y., Asada, K. (1981): Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology* 22: 867-880.
- Oxford Biomedical Research (2001): Glutathione reductase assay kit. Spectrophotometric Assay for Glutathione Reductase. Product No. FR 19: <http://www.oxfordbiomed.com/objects/catalog/product/extras/FR19.pdf> (Accessed March 2010)
- Paradičković, N., Vinković, T., Radman, D. (2008): Utjecaj biostimulatora na klijavost sjemena cvjetnih vrsta. *Sjemenarstvo* 25(1): 25-33.
- Paradičković, N., Zeljković, S., Đurić, G., Vinković, T., Mustapić-Karlić, J., Kanižai, G., Iljić, D. (2009): Rast i razvoj kadife (*Tagetes erecta* L.) pod utjecajem volumena supstrata i tretmana biostimulatorom. *Zbornik radova 44. hrvatskog i 4. međunarodnog simpozija agronoma. Lončarić, Z.; Marić, S. (ur.). Osijek: Sveučilište J. J. Strossmayera, Poljoprivredni fakultet u Osijeku, Opatija, Hrvatska, 2009: 786-790.*
- Redžepović, S., Čolo, J., Pecina, M., Duraković, L. (2008): Utjecaj biostimulatora rasta i fungicida za tretiranje sjemena soje na učinkovitost simbioze fiksacije dušika. *Sjemenarstvo* 24(3-4): 169-176.
- Roussos, P.A., Denaxa, N.K., Damvakaris, T. (2009): Strawberry fruit quality attributes after application of plant growth stimulating compounds. *Scientia Horticulturae* 119: 138-146.
- Schachtman, D.P., Shin, R. (2007): Nutrient Sensing and Signaling: NPKS. *Annual Review of Plant Biology* 58: 47-69.
- Shin, R., Schachtman, D.P. (2004): Hydrogen peroxide mediates plant root response to nutrient deprivation. *Proceedings of the National Academy of Sciences of the United States of America* 101: 8827-8832.
- Siegel, B.Z., Galston, W. (1967): The peroxidase of *Pisum sativum*. *Plant Physiology* 42: 221-226.

20. Takeda, F. (2000): Out-of-Season greenhouse strawberry production in soilless substrate. *Advances in Strawberry Research* 18: 4-15.
21. Turhan, E., Gulen, H., Eris, A. (2008): The activity of anti-oxidative enzymes in three strawberry cultivars related to salt-stress tolerance. *Acta Physiologiae Plantarum* 30: 201-208.
22. Vernieri, P., Malorgio, F., Tognoni, F. (2002): Use of biostimulants in production of vegetable seedlings. *Colture-Protette (Italy)*, 31(1): 75-79.
23. Zhang, X., Schmidt, R.E. (1999): Biostimulating turf-grasses. *Grounds Maintenance*, November, 1999: http://grounds-mag.com/mag/grounds_maintenance_biostimulating_turfgrasses/ (Accessed December 2009)

ANTIOKSIDATIVNI ODGOVOR U LISTU JAGODA (*Fragaria x ananassa* Duch) NA TRETMANE BIOSTIMULATORIMA UZ REDUCIRANU GNOJIDBU DUŠIKOM I KALIJEM

SAŽETAK

Sorta jagoda Elsanta uzgajana je u tresetnom supstratu u plasteniku. Puna doza i 50% smanjena doza dušika i kalija primjenjene su tijekom proljetnoga plodonošenja, uz primjenu biostimulatora Viva® i Megafol® i njihove kombinacije. Specifične aktivnosti gvajakol peroksidaze (GPXs; EC 1.11.1.7), katalaze (CATs; EC 1.11.1.6), askorbat peroksidaze (APXs; EC 1.11.1.11) i glutation reduktaze (GRs; EC 1.6.4.2) u listu jagode bile su značajno više poslije tretiranja biostimulatorima i reduciranom gnojdbom. Najjača povezanost uočena u ovom istraživanju, bila je između enzima askorbat-glutation ciklusa (APXs i GRs), koji su bili u pozitivnoj korelaciji s masom svježega lista (TLFM). Najveća masa svježega lista (TLFM) uočena je na biljkama tretiranim Megafolom®.

Ključne riječi: antioksidativna aktivnost, askorbat peroksidaza, biostimulatori, katalaza, gnojdba, *Fragaria x ananassa* Duch, glutation reduktaza, gvajakol peroksidaza, list, jagoda

(Received on 16 March 2010; accepted on 20 May 2010 - *Primljeno 16. ožujka 2010.; prihvaćeno 20. svibnja 2010.*)