

# Candidate Genes Associated with Economically Important Traits of Sheep - A Review

---

**Gavran, Mirna; Antunović, Zvonko; Gantner, Vesna**

*Source / Izvornik:* **Agriculturae Conspectus Scientificus, 2021, 86, 195 - 201**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

*Permanent link / Trajna poveznica:* <https://um.nsk.hr/um:nbn:hr:151:833879>

*Rights / Prava:* [In copyright](#)/[Zaštićeno autorskim pravom.](#)

*Download date / Datum preuzimanja:* **2025-03-16**



Sveučilište Josipa Jurja  
Strossmayera u Osijeku

**Fakultet  
agrobiotehničkih  
znanosti Osijek**

*Repository / Repozitorij:*

[Repository of the Faculty of Agrobiotechnical  
Sciences Osijek - Repository of the Faculty of  
Agrobiotechnical Sciences Osijek](#)



# Candidate Genes Associated with Economically Important Traits of Sheep - A Review

---

Mirna GAVRAN  
Zvonko ANTUNOVIĆ  
Vesna GANTNER (✉)

## Summary

---

Sheep give meat, milk, wool to the human population and this makes them a significant part of the world livestock industry. The basis of meat, milk and wool production is reproduction, respectively the success of reproduction. The result of reproduction is under the direct influence of genotypes and external factors and their interaction. The most important sheep product is meat and it is the main reason why sheep are raised in most countries. Milk is an important product of sheep, it is the basic food of young lambs and it plays an important role in the diet of the population. Wool makes sheep unique and therefore they are recognizable at first sight, in other words, different from other types of animals. Wool is a characteristic product of different and very variable economic importance, in some countries it is the most important product of sheep which provides the largest part of income. Many genes correlated with diverse economically significant phenotypic characteristics and their roles have been reported in domestic animals to date. In animal breeding, the identification of candidate genes correlated with economically important traits of sheep is crucial. The detailed information of the classified genes connected with significant traits helps to improve production capacity and productivity of sheep. Moreover, in this way the sheep processing industry can be increased, and thus the profitability and benefit of sheep breeders. The aim of this review is to provide complete information on the identified candidate genes associated with important production traits of sheep.

## Key words

---

sheep, meat, milk, wool, candidate genes

Faculty of Agrobiotechnical Sciences Osijek, Vladimira Preloga 1, 31000, Osijek, Croatia

✉ Corresponding author: [vgantner@fazos.hr](mailto:vgantner@fazos.hr)

Received: November 5, 2020 | Accepted: September 23, 2021

## Introduction

10,000 years ago, in the area of the Fertile Crescent, one of the earliest domesticated animal species were sheep. After domestication, a wide diversity of sheep breeds developed during the spreading to other regions (Larson et al., 2014; Lv et al., 2015; Gorkhali et al., 2016; Xu and Li, 2017). Sheep give meat, milk, wool to the human population and this makes them a significant part of the world livestock industry (Safari et al., 2005; Xu and Li, 2017). According to Martínez et al. (2016), the genomic region accountable for the variation of a quantitative characteristic is QTL. Also there is a genome-wide association study (GWAS). According to Zhang et al. (2012), the genome-wide association study (GWAS) is essential to identify candidate genes for many quantitative traits. To date, in domestic animals, many genes correlated with diverse economically significant phenotypic characteristics, and their roles have been reported. The aim of this review is to provide complete information on the identified candidate genes associated with important production traits of sheep.

## Methodology

During literature analysis, about one hundred references were screened and the 69 were selected as the most appropriate for the realisation of the paper goal. In the search, key search terms were: candidate genes; milk, wool, and meat sheep production; economically important traits in sheep. Relevance and recentness were set as selection criteria for inclusion or exclusion of studies.

## Genes Correlated with Fertility Traits

The basis of meat, milk and wool production is reproduction, respectively the success of reproduction. The result of reproduction is under the direct influence of genotypes and external factors and their interaction (Mioč et al., 2007). Fertility traits like ovulation rate, litter size, total number of lambs born, age at first lambing, age at first puberty and stillbirth have a great influence on the economy of the sheep industry. Ovulation rate and litter size are the most significant characteristics and also have high economic worth (Chen et al., 2015; Gebreselassie et al., 2020). Various genes control fertility as well as fecundity features with diverse influences on sheep (Drouilhet et al., 2009; Gebreselassie et al., 2020). Genes associated with ovulation rate, litter size, and sterility of sheep breeds have been identified by the GWAS study. These are the genes: growth differentiation factor 9 (GDF9), bone morphogenetic protein receptor IB (BMPRII) and bone morphogenetic protein 15 (BMP15) (Galloway et al., 2000; Souza et al., 2001; Hanrahan et al., 2004; Gebreselassie et al., 2020). Ovulation level and litter size are affected by BMPRII, while BMP15 along with GDF9 play an important role in follicle formation. The GDF9 gene is a protein belonging to the TGF- $\beta$  family and it additionally controls the development of the ovarian follicle. On the other hand, granulosa cells, theca cells and oocytes are affected by BMP15 (Davis et al., 1982; Yan et al., 2001; McNatty et al., 2005; Gebreselassie et al., 2020). In addition to fertility traits, there are also reproductive performance traits. Studies have shown genetic variants associated with reproductive abilities in some sheep breeds. A gene correlated with sensitivity to sheep lentivirus has been identified by GWAS study and its role is to

reduce sensitivity to lentivirus (Gebreselassie et al., 2020). Genes, fertility and reproductive traits and sheep breeds are shown in Table 1.

**Table 1.** Candidate genes responsible for fertility traits in sheep (Gebreselassie et al., 2020)

Genes	Chromosome	Traits	Sheep Breed
<i>PRLR</i>	16	RP	Herdwick & RFD
<i>TMEM154</i>	17	ILV	Herdwick & RFD
<i>CCNB2</i>	7	OD	GMM, CMF & AWD
<i>SLC8A3</i>	7	OD	GMM, CMF & AWD
<i>GDF9</i>	5	ORS	Cambridge&Belclare
<i>BMP15</i>	X	ORS	Cambridge&Belclare
<i>BMPRII</i>	6	ORS	Lacaune Sheep
<i>BMPRII</i>	7	LS	Han sheep
<i>BMP15</i>	X	LS	Han sheep

\*RP - reproductive performance; ILV - infection to the lentivirus; OD - oocyte development; ORS - ovulation rate & sterility; LS - litter size; GMM - German mutton merino; AWD - African white dorper; RFD - Rough Fells & Dalesbred; CMF - Chinese Mongolian fat-tailed

## Genes Correlated with Meat Production Traits

The most important sheep product is meat and it is the reason why sheep are raised in most countries. Lamb is much tastier and more sought-after, therefore fattening of adult sheep is almost neglected. For example, in New Zealand, of the total annual number of slaughtered sheep of different categories, 80% are lambs (Mioč et al., 2007). Even though mutton accounts for just 6% of world meat consumption, in regions of North Africa, the Middle East, India, and parts of Europe it is the main meat. The largest consumer of lamb and the first importer of lamb is the European Union; 99% of imported lamb comes from Australia and New Zealand (FAO, 2013). The best known and most thoroughly studied QTLs in sheep have been associated with muscle hypertrophy. The most described callipyge phenotype is connected with postnatal hypertrophy of muscle tissue with the concurrent decrease of carcass fatness (Cockett et al., 1996; Cockett et al., 2005; Knapik et al., 2017). The major gene that significantly influences carcass composition is Callipyge gene. In some sheep populations, a mutated variant of the normal (*clpg*) gene is located on the distal part of chromosome 18. Lower daily feed intake (Jackson et al., 1997; Knapik et al., 2017), better food efficiency and better slaughter quality are characteristics of Callipyge lambs, compared to lambs without this specified gene. This gene was originally identified in the herd of American Dorset and is inherited dominantly. Increased hardness of some muscles (Koochmarai et al., 1995; Shackelford et al., 1997; Knapik et al., 2017) is one of the worst effects of the Callipyge gene (Mioč, 2020). The Callipyge gene was first observed in the United States in 1983. Animals with the Callipyge phenotype have an increased muscle mass of 30%, particularly in the hindquarter. When

lambs of both sexes are about a month old, this phenomenon occurs and results in a higher proportion and a larger diameter of fast-twitching muscle fibers (Jackson et al., 1997; Reflexions, 2016; Knapik et al., 2017). Callipyge phenomenon is interesting to geneticists because of one very peculiar particularity: unique mode of heredity transmission (Reflexions, 2016). According to Cockett et al., 1996 and Knapik et al., 2017, only in heterozygous offspring that inherit the CLPG mutation from their fathers is the callipyge mutation phenotypically displayed. This unique mode of inheritance is called polar overdominance.

Another gene related to meat tenderness and growth performance characteristics in sheep is Calpain (CAPN) gene (Mahrous et al., 2016; Arora et al., 2019; Kumar et al. 2015). In Chall and Zel sheep breeds, meat quality and fatty acid composition are influenced by genetic variants that have been recognized in the calpastatin (CAST) gene (Aali et al., 2017; Gebreselassie et al., 2020). Also a gene that affects tenderness and rib eye (loin) muscling is the Carwell gene (Nicoll et al., 1998; Jopson et al., 2001; Warner et al., 2010). Myostatin actively inhibits the development of skeletal muscle (Bellings et al., 2005), and is also identified by reducing in total lean meat and increasing of the proportion of loin meat (Hickford et al., 2010; Gebreselassie et al., 2020). Genes for meat production traits and sheep breeds are shown in Table 2.

### Genes Correlated with Milk Traits

Milk is an important product of sheep, it is the basic food of young lambs and it plays an important role in the diet of the entire population. Sheep's milk is most often consumed in the form of cheese (Mioč et al., 2007). An important part in the agricultural economies of many countries, especially in the Mediterranean and Middle East regions, is covered by dairy sheep industry (Selvaggi et al., 2014; Suárez-Vega et al., 2016). Prolactin (PRL), leptin (LEP), insulin-like growth factor hormone 1 (IGF1), diacylglycerol O-acyltransferase 1 (DGAT1), signal transducer and activator of transcription 5 (STAT5), alpha ( $\alpha$ )s2-casein (CSN1S2), growth hormone receptor (GHR), and growth hormone-releasing hormone receptor (GHRHR) are very important candidate genes connected with milk and growth traits (Moioli et al., 2007; Orford et al., 2010; Staiger et al., 2010). Prolactin is crucial for lactation and has a significant function in milk production (Knight, 2001), while leptin has a significant part in animal growth and metabolism (Choudhary et al., 2005). IGF1 gene affects the litter size and the persistence of lactation (Scatà et al., 2010; He et al., 2012; Luridiana et al., 2020). DGAT1 is perhaps connected with variation in milk fat content (Kumar et al., 2016), and has a significant part in triacylglycerol biosynthesis, likewise in milk and growth traits (Chen et al., 2002; Abousoliman et al., 2020).

**Table 2.** Candidate genes responsible for meat production traits in sheep (Knapik et al., 2017)

Gene symbol	Gene name	Traits	Breed
<i>CLPG</i>	Callipyge	MH, CF; Impact on FE and DFI	American Dorset
<i>Carwell locus</i>	The rib-eye muscling locus (REM), Carwell	MH – <i>longissimus dorsi</i> muscle; increases the rib-eye area – about 11% and weight of the <i>longissimus dorsi</i> muscle – about 7%	Australian Poll Dorset
<i>MSTN (GDF8)</i>	Myostatin	Increases muscularity; Effect MC and FC in lamb carcasses	Belgian Texel Texel sheep
<i>CAST</i>	Calpastatin	LWG, CW, <i>longissimus dorsi</i> shear force; Impact on daily gain birth to weaning	Dorset Down, Dorset Down x Coopworth, Kurdi sheep, Chall Iranian sheep, Zel Iranian sheep
<i>FABP4</i>	Fatty acid binding protein 4	MT, MS, IFC in <i>longissimus dorsi</i>	Small-Tailed Han sheep, Tan sheep, Inner Mongolia sheep
<i>DGAT1</i>	Diacylglycerol acyltransferase 1	MT, IMF content and MS	Small-Tailed Han sheep, Tan sheep, Inner Mongolia sheep
<i>LEP</i>	Leptin	Increase of fat-tail percent, total body fat weight	Iranian sheep breeds (Shal, Zandi, Zel)
<i>TNNC2</i>	Fast skeletal muscle troponin C	MT, MS pH and CW	Small-Tailed Han sheep, Tan sheep, Inner Mongolia sheep

\*MH – muscle hypertrophy; CF – carcass fatness; MC – meat content; FC – fat content; MT – meat tenderness; MS – marbling score; CW – carcass weight; FE – feed efficiency; DFI – daily feed intake; LWG – live weight gain; IFC – intramuscular fat content

STAT5 represents central role in signal transduction from prolactin to milk protein genes (Bole-Feysot et al., 1998; Abousoliman et al., 2020). Caseins represent approximately 80% of proteins in ruminant milk (Haug et al., 2007; Park et al., 2007; Balthazar et al., 2017; Abousoliman et al., 2020). Casein is composed of 4 fractions:  $\alpha$ S1-casein (6.7%),  $\alpha$ S2-casein (22.8%),  $\beta$ -casein (61.6%), and  $\kappa$ -casein (8.9%) (Selvaggi et al., 2014; Balthazar et al., 2017). The CN gene family, along with CSN1S1 and CSN1S2, is determined on bovine chromosome 6, which is correlated with milk components and milk quality (Boettcher et al., 2004; Kucerova et al., 2006; Caroli et al., 2009; Ardicli et al., 2018). A meaningful impact of the CSN1S2 locus was detected on milk protein content (Ardicli et al., 2018). GH and GHRH are important candidate genes because they control body growth and are correlated with growth success, body size, and meat quality in cattle (Waters et al., 2001; Di Stasio et al., 2005; Garrett et al., 2008; Reardon et al., 2010; Abousoliman et al., 2020). It is noticed in Turkish sheep breed that POU class 1 homeobox 1 (POU1F1) gene influences milk production (Ozmen et al., 2014; Gebreselassie et al., 2020) Furthermore, it has a major function in controlling the function of growth hormone as well as prolactin genes (Lan et al., 2009; Gebreselassie et al., 2020). Genes associated with milk traits are recognized in other sheep breeds. These genes are palmdelphin (PALMD), ring finger protein 145 (RFP145) and alpha-lactalbumin (LALBA) (García-Gómez et al., 2012; Moiola et al., 2013; Gebreselassie et al., 2020). Genes for milk and growth traits are presented in Table 3.

### Genes Correlated with Wool Traits

Wool makes sheep unique and therefore they are recognizable at first sight, in other words, different from other types of animals. Wool is a characteristic product of different and very variable economic importance, in some countries it is the most important product of sheep which provides the largest part of income. In Australia, 3/4 of sheep are raised for wool. Australia produces about a quarter of the world's unwashed wool, while Europe accounts for about 12% of total annual world wool production. (Mioč et al., 2006; Mioč et al., 2007). Fiber diameter, staple length, fineness dispersion and crimp are significant wool characteristics that define the quality of wool (Wang et al., 2014; Gebreselassie et al., 2020). Also, the focus is on fiber diameter coefficient of variation, shedding, color and hair follicles (Fu et al., 2016; Xu and Li, 2017). Genes affirmed to be correlated with sheep wool are Keratin-Associated Protein (KAP) and Follistatin (FST); in Chinese merino sheep, the FST gene, which is located on chromosome 16, was identified for wool fiber diameter and wool crimp traits (Ma et al., 2017). Wool quality is also associated with the keratin gene family (KAP6.1, KAP8.1, KAP8.2, KRTAP9-2, and KAP16.4) which controls the function of wool and the growth of hair follicles (Ma et al., 2017; Gebreselassie et al., 2020). A greatly valuable trait is coat color. Coat color helps to differentiate among breeds and varieties; for sheep breeds, it is a significant economic trait. According to Song and Li, 2017; Koseniuk et al., 2018; Gebreselassie et al., 2020, v-kit Hardy-Zuckerman 4

**Table 3.** Candidate genes responsible for milk and growth traits

Gene symbol	Gene name	Chr	Traits	Breed
<i>POU1F1</i>	POU class 1 homeobox 1	1	MY	Turkish sheep
<i>RFP145</i>	Ring finger protein 145	na	MY	Italian Altamurana
<i>LALBA</i>	Alpha -lactalbumin	3	MP	Spanish Churra
<i>GH</i>	Growth hormone	11	MT, BG Lower MY	All breeds Serrada Estrela
<i>GHRH</i>	Growth hormone-releasing hormone receptor	4	BG	All breeds
<i>PALMD</i>	Palmdelphin	na	MY	Italian Altamurana
<i>PRL</i>	Prolactin	20	MY, MC	All breeds
<i>LEP</i>	Leptin	5	AG, metabolism	All breeds
<i>IGF1</i>	Insulin-like growth factor hormone 1	3	LS, LP	All breeds
<i>DGAT1</i>	Diacylglycerol O-acyltransferase 1	9	MFC, MT, GT	All breeds
<i>STAT5</i>	Signal transducer and activator of transcription 5	11	Signal transduction from prolactin to milk protein	All breeds
<i>CSN1S2</i>	Alpha ( $\alpha$ )s2-casein	6	MPC	All breeds

\*Chr – Chromosome; AG – animal growth; LP – lactation persistence; LS – litter size; MY – milk yield; MP – milk production; MC – milk composition; MT- milk traits; GT – growth traits; na - not available; MFC - milk fat content; MPC – milk protein content; BG – body growth



**Table 4.** Candidate genes responsible for wool traits (Gebreselassie et al., 2020)

Gene symbol	Gene name	Chromosome	Trait	Breed
<i>KAP6.1</i>	Keratin associated protein 6.1	na	Wool	Merino (Xinjiang type)
<i>KAP8.1</i>	Keratin associated proteins 8.1	1	Wool	Merino (Xinjiang type)
<i>KAP8.2</i>	Keratin associated protein 8.2	na	Wool	Merino (Xinjiang type)
<i>KRTAP9-2</i>	Keratin associated protein 9-2	na	Wool	Merino (Xinjiang type)
<i>KAP16.4</i>	Keratin associated protein 16.4	na	Wool	Merino (Xinjiang type)
<i>FST</i>	Follistatin	16	WQ	CMS (Junken Type)
<i>MC1R</i>	Melanocortin 1 receptor	14	Color	MRA
<i>ASIP</i>	Agouti signaling protein	13	Color	Dubian & Privorianp

\*WQ - wool quality; na - not available; CMS - Chinese Merino Sheep; MRA - Manchega & Rasa Aragonesa

feline sarcoma viral oncogene homologue (KIT), agouti signaling protein (ASIP), microphthalmia-associated transcription factor (MITF) and the melanocortin 1 receptor (MC1R), tyrosinase-related protein 1 (TYRP1) are candidate genes responsible for coat color. Genes for wool traits are shown in Table 4.

## Conclusion

In animal breeding, the identification of candidate genes correlated with economically important traits of sheep is crucial. Many genes correlated with various phenotypic characteristics have been recorded in sheep. Certain stated genes influence more than one trait. The detailed information of the classified genes connected with significant traits helps to improve production capacity and productivity of sheep. Moreover, in this way the sheep processing industry can be increased, and thus the profitability and benefit of sheep breeders. In order to advance the effectiveness of sheep production, further research work in this field is needed.

## References

- Aali M., Moradi-Shahrbabak H., Moradi-Shahrbabak M., Sadeghi M., Yousefi A. R. (2017). Association of the Calpastatin Genotypes, Haplotypes, and SNPs with Meat Quality and Fatty Acid Composition in Two Iranian Fat- and Thin-Tailed Sheep Breeds. *Small Rumin Res* 149: 40–51. doi: 10.1016/j.smallrumres.2016.12.026
- Abousoliman I., Reyer H., Oster M., Muráni E., Mourad M., Abdel-Salam Rashed M., Mohamed I., Wimmers K. (2020). Analysis of Candidate Genes for Growth and Milk Performance Traits in the Egyptian Barki Sheep. *Animals* 10 (2): 197. doi: 10.3390/ani10020197.
- Ardicli S., Soyudal B., Samli H., Dincel D., Balci, F. (2018). Effect of STAT1, OLR1, CSN1S1, CSN1S2, and DGAT1 Genes on Milk Yield and Composition Traits of Holstein Breed. *Revista Brasileira de Zootecnia* 47: 1-9. doi: 10.1590/rbz4720170247.
- Arora R., Yadav H. S., Yadav D. K. (2019). Identification of Novel Single Nucleotide Polymorphisms in Candidate Genes for Mutton Quality in Indian Sheep. *Anim Mol Breed* 4 (1). doi: 10.5376/amb.2014.04.0001.
- Balthazar C. F., Pimentel T. C., Ferrão L. L., Almada C. N., Santillo A., Albenzio M., Mollakhalili N., Mortazavian A. M., Nascimento J. S., Silva M. C., Freitas M. Q., Sant'Ana A. S., Granato D., Cruz A. G.

- (2017). Sheep Milk: Physicochemical Characteristics and Relevance for Functional Food Development. *Compr Rev Food Sci Food Saf.* 16 (2): 247-262. doi: 10.1111/1541-4337.12250
- Bellinge R. H. S., Liberles D. A., Iaschi S. P. A., O'Brien P. A., Tay G. K. (2005). Myostatin and Its Implications on Animal Breeding: A Review. *Anim Genet* 36 (1): 1–6. doi: 10.1111/j.1365-2052.2004.01229.x
- Boettcher P., Caroli A., Stella A., Chessa S., Budelli E., Canavesi F., Ghiroldi S., Pagnacco G. (2004). Effects of Casein Haplotypes on Milk Production Traits in Italian Holstein and Brown Swiss Cattle. *J Dairy Sci* 87 (12): 4311-4317. doi: 10.3168/jds.S0022-0302(04)73576-6
- Bole-Feysot C., Goffin V., Edery M., Binart N., Kelly P.A. (1998). Prolactin (PRL) and Its Receptor: Actions Signal Transduction Pathways and Phenotypes Observed in PRL Receptor Knockout Mice. *Endocr Rev* 19 (3): 225–268. doi: 10.1210/edrv.19.3.0334
- Caroli A., Chessa S., Erhardt G. (2009). Invited Review: Milk Protein Polymorphisms in Cattle: Effect on Animal Breeding and Human Nutrition. *J Dairy Sci* 92: 5335-5352. doi: 10.3168/jds.2009-2461
- Chen H. C., Smith S. J., Ladha Z., Jensen D. R., Ferreira L. D., Pulawa L. K. (2002). Increased Insulin and Leptin Sensitivity in Mice Lacking Acyl CoA: diacylglycerol acyltransferase 1. *J Clin Invest* 109 (8): 1049–1055. doi: 10.1172/JCI14672.
- Chen H. Y., Shen H., Jia B., Zhang Y. S., Wang X. H., Zeng X. C. (2015). Differential Gene Expression in Ovaries of Qira Black Sheep and Hetian Sheep Using RNA-Seq Technique. *PLoS ONE* 10(3): e0120170. doi: 10.1371/journal.pone.0120170.
- Choudhary V., Kumar P., Bhattacharya T. K., Bhushan B., Sharma A. (2005). DNA Polymorphism of Leptin Gene in Bos Indicus and Bos Taurus Cattle. *Genet Mol Biol* 28 (4): 740–742. doi: 10.1590/S1415-47572005000500014
- Cockett N. E., Jackson S. P., Shay T. L., Farnir F., Berghmans S., Snowder G. D., Nielsen D., Georges M. (1996). Polar Overdominance at the Bovine *callipyge* locus. *Science* 273 (5272): 236-238. doi: 10.1126/science.273.5272.236
- Cockett N. E., Smit M. A., Bidwell C. A., Segers K., Hadfield T. L., Snowder G. D., Georges M., Charlier C. (2005). The *callipyge* Mutation and Other Genes What Affect Muscle Hypertrophy in Sheep. *Genet Sel Evol* 37 (Suppl 1): S65-81. doi: 10.1186/1297-9686-37-S1-S65
- Davis G., Montgomery G., Allison A., Kelly R., Bray A. (1982). Segregation of A Major Gene Influencing Fecundity in Progeny of Booroola Sheep. *N Z J Agric Res* 25 (4): 525–529. doi: 10.1080/00288233.1982.10425216.
- Di Stasio L., Destefanis G., Brugiapaglia A., Albera A., Rolando A. (2005). Polymorphism of the GHR Gene in Cattle and Relationships with Meat Production and Quality. *Anim Genet* 36 (2): 138-40. doi: 10.1111/j.1365-2052.2005.01244.x

- Drouilhet L., Lecerf F., Bodin L., Fabre S., Mulsant, P. (2009). Fine Mapping of the Fecl Locus Influencing Prolificacy in Lacaune Sheep. *Anim Genet* 40: 804–812.
- Food and Agriculture Organization, FAO (2013). FAOSTAT. DATA. Food Supply - Livestock and Fish Primary Equivalent. Available at: <http://www.fao.org/faostat/en/#data/CL> [Accessed 21 10. 2020].
- Fu S. Y., Qi Y. X., He X. L., Da L., Wang B., Rigele T. E., Wu J. H., Yang D., Liu Y. B., Zhang W. G. (2016). Transcriptome Analysis Reveals Skin Lipid Metabolism Related to Wool Diameter in Sheep. *BioRxiv*. doi: 10.1101/051359
- Galloway S. M., McNatty K. P., Cambridge L. M., Laitinen M. P., Juengel J. L., Jokiranta T. S., McLaren R. J., Luiro K., Dodds K. G., Montgomery, G. W. (2000). Mutations in An Oocyte-Derived Growth Factor Gene (BMP15) Cause Increased Ovulation Rate and Infertility in A Dosage-Sensitive Manner. *Nat Genet* 25 (3): 279–83. doi: 10.1038/77033
- Garrett A., Rincon G., Medrano J., Elzo M., Silver G., Thomas M. (2008). Promoter Region of the Bovine Growth Hormone Receptor Gene: Single Nucleotide Polymorphism Discovery in Cattle and Association with Performance in Brangus Bulls. *J Anim Sci* 86 (12): 3315–23. doi: 10.2527/jas.2008-0990
- Gebreselassie G., Berihulay H., Jiang L., Ma, Y. (2020). Review on Genomic Regions and Candidate Genes Associated with Economically Important Production and Reproduction Traits in Sheep (*Ovis aries*). *Animals* 10 (1): 33. doi: 10.3390/ani10010033
- Gorkhali N. A., Dong K. Z., Yang M., Song S., Kader A., Shrestha B. S., He X. H., Zhao Q. J., Pu Y. B., Li X. C., Kijas J., Guan W. J., Han J. L., Jiang L., Ma Y. H. (2016). Genomic Analysis Identified a Potential Novel Molecular Mechanism for High-Altitude Adaptation in Sheep at the Himalayas. *Sci Rep* 6 (1): 29963. doi: 10.1038/srep29963
- Hanrahan J. P., Gregan S. M., Mulsant P., Mullen M., Davis G. H., Powell R., Galloway S. M. (2004). Mutations in the Genes for Oocyte-Derived Growth Factors GDF9 and BMP15 are Associated with Both Increased Ovulation Rate and Sterility in Cambridge and Belclare Sheep (*Ovis aries*). *Biol Reprod* 70 (4): 900–909. doi: 10.1095/biolreprod.103.023093
- Haug A., Hostmark A. T., Harstad O. M. (2007). Bovine Milk in Human Nutrition—A Review. *Lipids Health Dis* 6(1): 25. doi: 10.1186/1476-511X-6-25
- He J. N., Zhang B. Y., Chu M. X., Wang P. Q., Feng T., Cao G. L., Di R., Fang L., Huang D. W., Tang Q. Q., Li N. (2012). Polymorphism of Insulin-like Growth Factor 1 Gene and Its Association with Litter Size in Small Tail Han Sheep. *Mol Biol Rep* 39: 9801–9807. doi: 10.1007/s11033-012-1846-y
- Hickford J., Forrest R., Zhou H., Fang Q., Han J., Frampton C., Horrell, A. (2010). Polymorphisms in the Ovine Myostatin Gene (MSTN) and Their Association with Growth and Carcass Traits in New Zealand Romney Sheep. *Anim Genet* 41 (1): 64–72. doi: 10.1111/j.1365-2052.2009.01965.x
- Jackson S. P., Green R. D., Miller M. F. (1997). Phenotypic Characterization of Rambouillet Sheep Expressing the Callipyge Gene: I. Inheritance of the Condition and Production Characteristics. *J Anim Sci* 75 (1): 14–18. doi: 10.2527/1997.75114x
- Jopson N., Nicoll G. B., Stevenson-Barry J. M., Duncan S., Greer G. J., Bain W., Gerard E., Glass B. C., Broad T. E., McEwan J. C. (2001). Mode of Inheritance and Effects on Meat Quality of the Rib-Eye Muscling (REM) QTL in Sheep. *Proceedings of the Association for the Advancement of Animal Breeding and Genetics, Queenstown, New Zealand* 14, pp. 111–114. doi: 10.13140/2.1.3002.8167.
- Knapik J., Ropka-Molik K., Pieszka M. (2017). Genetic and Nutritional Factors Determining the Production and Quality of Sheep Meat – A Review. *Ann Anim Sci* 17 (1): 23–40. doi: 10.1515/aoas-2016-0036
- Knight C. H. (2001). Overview of Prolactin's Role in Farm Animal Lactation. *Livest Prod Sci* 70 (1): 87–93. doi: 10.1016/S0301-6226(01)00200-7
- Koohmaraie M., Shackelford S.D., Wheeler T.L., Lonergan S.M., Doumit M.E. (1995). A Muscle Hypertrophy Condition in Lamb (*callipyge*): Characterization of Effects on Muscle Growth and Meat Quality Traits. *J Anim Sci* 73 (12): 3596–607. doi: 10.2527/1995.73123596x
- Koseniuk A., Ropka-Molik K., Rubiś D., Smolucha G. (2018). Genetic Background of Coat Colour in Sheep. *Arch Anim Breed* 61 (2): 173–178. doi: 10.5194/aab-61-173-2018
- Kucerova J., Matejcek A., Jandurova O., Sorensen P., Nemcova E., Stipkova M., Kott T., Bouska J., Frelich J. (2006). Milk Protein Genes CSN1S1, CSN2, CSN3, LGB and Their Relation to Genetic Values of Milk Production Parameters in Czech Fleckvieh. *Czech J Anim Sci* 51: 241–247. doi: 10.17221/3935-CJAS
- Kumar A., Chhokar V., Kumar R., Aggarwal H., Beniwa V. (2016). Short Communication Molecular Characterization of Acyl CoA: Diacylglycerol O-acyltransferase 1 (DGAT1) in Sheep and its Comparison with Other Ruminants. *Am J Biochem Mol Biol* 6 (2): 67–71. doi: 10.3923/ajbmb.2016.67.71.
- Kumar N., Jayashankar M., Ramakrishnappa N., Nagaraja C., Fairoze N., Satyanarayana K. (2015). Genetic Polymorphism of Ovine Calpain Gene in Bandur Sheep. *Int J Sci Environ Technol* 4 (3): 804–812
- Lan X., Li M., Chen H., Zhang L., Jing Y., Wei T., Ren G., Wang X., Fang X., Zhang, C. (2009). Analysis of Caprine Pituitary Specific Transcription Factor-1 Gene Polymorphism in Indigenous Chinese Goats. *Mol Biol Rep* 36 (4): 705–709. doi: 10.1007/s11033-008-9232-5
- Larson G., Piperno D. R., Allaby R. G., Purugganan M. D., Andersson L., Arroyo-Kalin M., Barton L., Climer Vigueira C., Denham T., Dobney K., Doust A. N., Gepts P., Gilbert M. T. P., Gremillion K. J., Lucas L., Lukens L., Marshall F. B., Olsen K. M., Pires J. C., Richerson P. J., Rubio de Casas R., Sanjurjo O. I., Thomas M. G., Fuller D. Q. (2014). Current Perspectives and the Future of Domestication Studies. *Proceedings of the National Academy of Sciences of the United States of America*, 111(17), pp. 6139–6146.
- Luridiana S., Mura M. C., Di Stefano M. V., Pulinas L., Cosso G., Nehme M., Carcangiu V. (2020). Polymorphism of Insulin-like Growth Factor 1 Gene and Its Relationship with Reproductive Performances and Milk Yield in Sarda Dairy Sheep. *Vet Anim Sci* 9: 100084. doi: 10.1016/j.vas.2019.100084
- Lv F. H., Peng W. F., Yang J., Zhao Y. X., Li W. R., Liu M. J., Ma Y. H., Zhao Q. J., Yang G. L., Wang F., Li J. Q., Liu Y. G., Shen Z. Q., Zhao S. G., Hehua E. E., Gorkhali N. A., Farhad Vahidi S. M., Muladno M., Naqvi A. N., Tabell J., Iso-Touru T., Bruford M. W., Kantanen J., Han J. L., Li M. H. (2015). Mitogenomic Meta-Analysis Identifies Two Phases of Migration in the History of Eastern Eurasian Sheep. *Mol Biol Evol* 32 (10): 2515–2533
- Ma G. W., Chu Y. K., Zhang W. J., Qin F. Y., Xu S. S., Yang H., Rong E. G., Du Z. Q., Wang, S. Z., Li H. (2017). Polymorphisms of FST Gene and Their Association with Wool Quality Traits in Chinese Merino Sheep. *PLoS ONE* 12 (4): e0174868. doi: 10.1371/journal.pone.0174868
- Mahrous K., Hassanane M., Shafey H., Mordy M.A., Rushdi, H. (2016). Association between Single Nucleotide Polymorphism in Ovine Calpain Gene and Growth Performance in Three Egyptian Sheep Breeds. *J Genet Eng Biotechnol* 14 (2): 233–240. doi: 10.1016/j.jgeb.2016.09.003
- Martínez P., Robledo D., Rodríguez-Ramilo S. T., Hermida M., Taboada X., Pereiro P., Rubiolo J.A., Ribas L., Gómez-Tato A., Álvarez-Dios J. A., Piferrer F., Novoa B., Figueras A., Pardo B.G., Fernández J., Viñas A., Bouza, C. (2016). 6 - Turbot (*Scophthalmus maximus*) Genomic Resources: Application for Boosting Aquaculture Production, *Genomics in Aquaculture*, pp. 131–163. doi: 10.1016/B978-0-12-801418-9.00006-8
- McNatty K. P., Galloway S. M., Wilson T., Smith P., Hudson N. L., O'Connell A., Bibby A. H., Heath D. A., Davis G. H., Hanrahan J. P. (2005). Physiological Effects of Major Genes Affecting Ovulation Rate in Sheep. *Genet Sel Evol* 37 (Suppl. 1): S109–S123. doi: 10.1186/1297-9686-37-S1-S25
- Mioč B. (2020). Selekcija ovaca za meso. (Hrvatski savez uzgajivača ovaca i koza) Available at: <http://www.ovce-koze.hr/ovcarstvo-kozarstvo/edukativni-kutak/selekcija-ovaca-za-meso> [Accessed 21 09. 2020]
- Mioč B., Pavić V., Sušić V. (2007). Ovcarstvo. Hrvatska mljekarska udruga, Zagreb. 424 pp.

- Mioč B., Sušić V., Pavić V., Barać Z., Prpić Z. (2006). Priprema ovaca za strižu, striža i postupci s vunom do transporta. *Stočarstvo* 60 (2): 129-141
- Moioli B., D'Andrea M., Pilla F. (2007). Candidate Genes Affecting Sheep and Goat Milk Quality. *Small Ruminant Res* 68 (1-2): 179-192. doi: 10.1016/j.smallrumres.2006.09.008
- Moioli B., Scatà M.C., Steri R., Napolitano F., Catillo G. (2013). Signatures of Selection Identify Loci Associated with Milk Yield in Sheep. *BMC Genet* 14: 76. doi: 10.1186/1471-2156-14-76
- Nicoll G. B., Burkin H. R., Broad T. E., Jopson N., Greer G. J., Bain W., Wright C. S., Dodds K. G., Fennessy P., McEwan J. C. (1998). Genetic Linkage of Microsatellite Markers to the Carwell Locus for Rib-Eye Muscling in Sheep. *Proceedings of the 6th World Congress of Genetics Applied to Livestock Production*, 26, pp. 529-532. doi: 10.13140/2.1.1379.9684
- Orford M., Tzamaloukas O., Papachristoforou C., Miltiadou D. (2010). Technical Note: A Simplified PCR Based Assay for the Characterization of Two Prolactin Variants that Affect Milk Traits in Sheep Breeds. *J Dairy Sci* 93 (12): 5996-5999. doi: 10.3168/jds.2010-3569
- Ozmen O., Kul S., Unal E.O. (2014). Polymorphism of Sheep POU1F1 Gene Exon 6 and 30UTR Region and Their Association with Milk Production Traits. *Iran J Vet Res* 15 (4): 331-335
- Park Y.W., Juárez M., Ramos M., Haenlein G.F.W. (2007). Physico-Chemical Characteristics of Goat and Sheep Milk. *Small Rumin Res* 68 (1-2): 88-113. doi: 10.1016/j.smallrumres.2006.09.013
- Reardon W., Mullen A., Sweeney T., Hamill R. (2010). Association of Polymorphisms in Candidate Genes with Colour, Water-Holding Capacity and Composition Traits in Bovine *M. longissimus* and *M. semimembranosus*. *Meat Sci* 86 (2): 270-5. doi: 10.1016/j.meatsci.2010.04.013.
- Reflexions (2016). The Callipyge Phenomenon, a Genetic Curiosity. The University of Liège. Available at: [http://www.reflexions.uliege.be/cms/c\\_423320/en/the-callipyge-phenomenon-a-genetic-curiosity](http://www.reflexions.uliege.be/cms/c_423320/en/the-callipyge-phenomenon-a-genetic-curiosity) [Accessed 28 09. 2020 ].
- Safari E., Fogarty N. M., Gilmour A. R. (2005). A Review of Genetic Parameter Estimates for Wool, Growth, Meat and Reproduction Traits in Sheep. *Livest Prod Sci* 92 (3): 271-289
- Scatà M. C., Catillo G., Annicchiarico G., De Matteis G., Napolitano F., Signorelli F., Moioli B. (2010). Investigation on Lactation Persistency and IGF-I Gene Polymorphisms in Dairy Sheep. *Small Rumin Res* 89, (1): 7-11. doi: 10.1016/j.smallrumres.2009.10.014
- Selvaggi M., Laudadio V., Dario C., Tufarelli V. (2014). Investigating the Genetic Polymorphism of Sheep Milk Proteins: A Useful Tool for Dairy Production. *J Sci Food Agric* 94 (15): 3090-3099. <http://dx.doi.org/10.1002/jsfa.6750>.
- Shackelford S. D., Wheeler T. L., Koohmaraie M. (1997). Effect of the Callipyge Phenotype and Cooking Method on Tenderness of Several Major Lamb Muscles. *J Anim Sci* 75 (8): 2100-2105. doi: 10.2527/1997.7582100x
- Souza C., MacDougall C., Campbell B., McNeilly A., Baird D. (2001). The Booroola (FecB) Phenotype Is Associated with a Mutation in the Bone Morphogenetic Receptor Type 1 B (BMPRI1B) Gene. *J Endocrinol* 169 (2): R1-6. doi: 10.1677/joe.0.169r001
- Staiger E. A., Thonney M. L., Buchanan J. W., Rogers E. R., Oltenacu P. A., Mateescu R. G. (2010). Effect of Prolactin  $\beta$ -Lactoglobulin and  $\kappa$ -Casein Genotype on Milk Yield in East Friesian Sheep. *J Dairy Sci* 93: 1736-1742. doi: 10.3168/jds.2009-2630
- Suárez-Vega A., Gutiérrez-Gil B., Arranz J. J. (2016). Transcriptome Expression Analysis of Candidate Milk Genes Affecting Cheese-Related Traits in 2 Sheep Breeds. *J Dairy Sci* 99: 6381-6390. doi: 10.3168/jds.2016-11048
- Wang Z., Zhang H., Yang H., Wang S., Rong E., Pei W., Li H., Wang, N. (2014). Genome-Wide Association Study for Wool Production Traits in a Chinese Merino Sheep Population. *PLoS ONE* 9 (9): e107101. doi: 10.1371/journal.pone.0107101
- Warner R. D., Greenwood P. L., Pethick D. W., Ferguson D. M. (2010). Review. Genetic and Environmental Effects on Meat Quality. *Meat Sci* 86 (1): 171-183. doi: 10.1016/j.meatsci.2010.04.042
- Waters S., McCabe M., Howard D., Giblin L., Magee D., MacHugh D., Berry D. (2001). Associations between Newly Discovered Polymorphisms in the Bos Taurus Growth Hormone Receptor Gene and Performance Traits in Holstein-Friesian Dairy Cattle. *Anim Genet* 42 (1): 39-49. doi: 10.1111/j.1365-2052.2010.02087.x
- Xu S. S., Li M. H. (2017). Recent Advances in Understanding Genetic Variants Associated with Economically Important Traits in Sheep (*Ovis aries*) Revealed by High-Throughput Screening Technologies. *Front Agr Sci Eng* 4 (3): 279-288. doi: 10.15302/J-FASE-2017151
- Yan C., Wang P., DeMayo J., DeMayo F. J., Elvin J. A., Carino C., Prasad S. V., Skinner S.S., Dunbar B.S., Dube, J.L. (2001). Synergistic Roles of Bone Morphogenetic Protein 15 and Growth Differentiation Factor 9 in Ovarian Function. *Mol Endocrinol* 15 (6): 854-866. doi: 10.1210/mend.15.6.0662
- Zhang H., Wang Z., Wang S., Li H. (2012). Progress of Genome Wide Association Study in Domestic Animals. *J Anim Sci Biotechnol* 3: 26. doi: 10.1186/2049-1891-3-26